Czechoslovak Mathematical Journal, 55 (130) (2005), 283–293

ON SIGNPOST SYSTEMS AND CONNECTED GRAPHS

Ladislav Nebeský, Praha

(Received May 20, 2002)

Abstract. By a *signpost system* we mean an ordered pair (W, P), where W is a finite nonempty set, $P \subseteq W \times W \times W$ and the following statements hold:

if $(u, v, w) \in P$, then $(v, u, u) \in P$ and $(v, u, w) \notin P$, for all $u, v, w \in W$;

if $u \neq v$, then there exists $r \in W$ such that $(u, r, v) \in P$, for all $u, v \in W$.

We say that a signpost system (W, P) is *smooth* if the folowing statement holds for all $u, v, x, y, z \in W$: if $(u, v, x), (u, v, z), (x, y, z) \in P$, then $(u, v, y) \in P$. We say thay a signpost system (W, P) is *simple* if the following statement holds for all $u, v, x, y \in W$: if $(u, v, x), (x, y, v) \in P$, then $(u, v, y), (x, y, u) \in P$.

By the underlying graph of a signpost system (W, P) we mean the graph G with $V(G) = W$ and such that the following statement holds for all distinct $u, v \in W$: u and v are adjacent in G if and only if $(u, v, v) \in P$. The main result of this paper is as follows: If G is a graph, then the following three statements are equivalent:

G is connected;

G is the underlying graph of a simple smooth signpost system;

G is the underlying graph of a smooth signpost system.

Keywords: connected graph, signpost system

MSC 2000: 05C38, 05C40, 05C12

By a graph we mean here a *finite* undirected graph with no multiple edges or loops. The letters $f - n$ will serve for denoting non-negative integers.

1. Ternary systems

By a *ternary system* we mean an ordered pair (W, P) such that W is a finite nonempty set and $P \subseteq W \times W \times W$. Obviously, if (W, P) is a ternary system, then P is a ternary relation in W.

Research was supported by Grant Agency of the Czech Republic, grant No. 401/01/0218.

Let $S = (W, P)$ be a ternary system. Consider $u_0, \ldots, u_i, v \in W$, where $i \geqslant 1$. If

$$(u_j, u_{j+1}, v) \in P \quad \text{for each } j,\ 0 \leqslant j \leqslant i-1,$$

we will write

$$u_0 \ldots u_i S v.$$

Thus, instead of $(u, v, w) \in P$, where $u, v, w \in W$, we write $uvSw$. Moreover, instead of $(x, y, z) \notin P$, where $x, y, z \in W$, we will write $\neg(xySz)$. Finally, we will write $V(S) = W$.

We say that a ternary system S is *smooth* if it satisfies the following axiom (SMO):

(SMO) if $uvSx,\ uvSz$ and $xySz,$ then $uvSy$ for all $u, v, x, y, z \in V(S)$.

Lemma 1. *Let S be a smooth ternary system and let $u_0, \ldots, u_i, x, y, z \in V(S)$, where $i \geqslant 1$. Assume that $u_0 \ldots u_i Sx$, $u_0 \ldots u_i Sz$ and $xySz$. Then $u_0 \ldots u_i Sy$.*

P r o o f. We proceeed by induction on i. The case when $i = 1$ follows immediately from (SMO). Let $i \geqslant 2$. Obviously, $u_0 \ldots u_{i-1} Sx$ and $u_0 \ldots u_{i-1} Sz$. By the induction hypothesis, $u_0 \ldots u_{i-1} Sy$. Recall that $u_{i-1} u_i Sx$, $u_{i-1} u_i Sz$ and $xySz$. As follows from (SMO), $u_{i-1} u_i Sy$. Thus $u_0 \ldots u_i Sy$, which completes the proof. □

Proposition 1. *Let S be a smooth ternary system, and let $u_0, \ldots, u_i, x_0, \ldots, x_j \in V(s)$, where $i, j \geqslant 1$. If*

(1) $$u_0 \ldots u_i S x_0, \quad u_0 \ldots u_i S z \quad \textit{and} \quad x_0 \ldots x_j S z,$$

then

(2) $$u_0 \ldots u_i S x_j.$$

P r o o f. Assume that (1) holds. We will prove that (2) holds. We proceed by induction on j. The case when $j = 1$ follows immediately from Lemma 1. Let $j \geqslant 2$. Obviously, $x_0 \ldots x_{j-1} Sz$. By the induction hypothesis, $u_0 \ldots u_i S x_{j-1}$. Since $u_0 \ldots u_i Sz$ and $x_{j-1} x_j Sz$, Lemma 1 implies that $u_0 \ldots u_i S x_j$, which completes the proof. □

We say that a ternary system S is *simple* if it satisfies the following axioms (SIM1) and (SIM2):

(SIM1) if $uvSx,\ xySv,$ then $uvSy$ for all $u, v, x, y \in V(S)$;

(SIM2) if $uvSx,\ xySv,$ then $xySu$ for all $u, v, x, y \in V(S)$.

The next lemma depends on a result proved in [4].

Lemma 2. *Let S be a simple ternary system, let $w_0, \dots, w_k$, $y, z \in V(S)$, where $k \geqslant 1$. Assume that $w_0 \dots w_k Sy$. If $zySw_0$, then $w_0 \dots w_k Sz$ and $zySw_k$. If $yzSw_k$, then $w_0 \dots w_k Sz$ and $yzSw_0$.*

P r o o f. The lemma immediately follows from Lemma 1 in [4]. □

Proposition 2. *Let S be a simple ternary system, and let $u_0, \dots u_i, x_0, \dots, x_j \in V(S)$, where $i, j \geqslant 1$. If*

$$u_i \dots u_0 Sx_0 \quad \text{and} \quad x_0 \dots x_j Su_0, \tag{3}$$

then

$$u_i \dots u_0 Sx_j \quad \text{and} \quad x_0 \dots x_j Su_i. \tag{4}$$

P r o o f. Put $h = i + j$. Obviously, $h \geqslant 2$. Let (3) hold. We will prove that (4) holds. We proceeed by induction on h. First, let $h = 2$. Then $i = 1 = j$. Clearly, (SIM1) and (SIM2) imply (4).

Let $h \geqslant 3$. If $i = 1$ or $j = 1$, then (4) immediatelly follows from Lemma 2. Assume that $i, j \geqslant 2$. Then $h \geqslant 4$. Since $u_i \dots u_0 Sx_0$ and $x_0 \dots x_{j-1} Su_0$, the induction hypothesis implies that $u_i \dots u_0 Sx_{j-1}$. Since $u_{i-1} \dots u_0 Sx_0$ and $x_0 \dots x_j Su_0$, the induction hypothesis implies that $x_0 \dots x_j Su_{i-1}$. It follows from (3) that $u_i u_{i-1} Sx_0$ and $x_{j-1} x_j Su_0$. Since $x_0 \dots x_j Su_{i-1}$ and $u_i u_{i-1} Sx_0$, Lemma 2 implies that $x_0 \dots x_j Su_i$. Since $u_i \dots u_0 Sx_{j-1}$ and $x_{j-1} x_j Su_0$, Lemma 2 implies that $u_i \dots u_0 Sx_j$. Hence (4) holds. □

2. Signpost systems and their underlying graphs

By a *signpost system* we mean a ternary system S satisfying the following axioms (SIG1), (SIG2) and (SIG3):

(SIG1) if $uvSw$, then $vuSu$ for all $u, v, w \in V(S)$,

(SIG2) if $uvSw$, then $\neg(vuSw)$ for all $u, v, w \in V(S)$,

(SIG3) if $u \neq v$, then there exists $r \in V(S)$ such that $urSv$ for all $u, v \in V(S)$.

Remark. If S is a signpost system, $u, v, w \in V(S)$ and $uvSw$, then the ordered triple (u, v, w) can be considered as a signpost showing a "direction" from u to w; this direction is determined by v.

The notion of a signpost system appeared in [2] but an implicit form of the idea of a signpost system can be found in [4] already. Our definition follows the definition in [6], which is slightly different from that in [2]. Note that (W, P) is a signpost system (in the sense of our definition) if and only if P is a signpost system on W in the sense of the definition in [2].

Proposition 3. *Let S be a signpost system. Then*

(5) $$\text{if } uvSw, \text{ then } uvSv \text{ for all } u, v, w \in V(S),$$

(6) $$uvSv \text{ if and only if } vuSu \text{ for all } u, v \in V(S)$$

and

(7) $$\text{if } uvSw, \text{ then } v \neq u \neq w \text{ for all } u, v, w \in V(S).$$

P r o o f. Obviously, (1) and (2) are implied by (SIG1). Combining (SIG1) and (SIG2) we get (3). □

Lemma 3. *Let S be a smooth signpost system. Then S satisfies the axiom* (SIM1).

P r o o f. Consider arbitrary $u, v, x, y \in V(S)$. Let $uvSx$, $xySv$. By (5), $uvSv$. If we put $z = v$ in (SMO), we get $uvSy$. □

Let S be a signpost system. By the *underlying* graph of S we mean the graph G defined as follows: $V(G) = V(S)$ and

$$u \text{ and } v \text{ are adjacent in } G \text{ if and only if } uvSv$$

for all $u, v \in V(S)$.

Let S be a signpost system. By a *confusing circuit* in S we mean an ordered pair

(8) $$(u_0 u_1 \dots u_k, v)$$

such that $u_0, u_1, \dots u_k, v \in V(S)$, $k \geqslant 1$, $u_0 \dots u_k S v$ and $u_k = u_0$.

Lemma 4. *Let S be a signpost system, and let* (8) *be a confusing circuit in S. Then $u_0 \neq v$ and $k \geqslant 3$.*

P r o o f. As follows from (7), $u_0 \neq v$ and $k \geqslant 2$. Let $k = 2$. Since $u_2 = u_0$, we get u_1u_0Sv and u_0u_1Sv, which contradicts (SIG2). Thus $k \geqslant 3$. □

Example. Let X, Y and Z be pairwise disjoint finite sets such that $|X| \geqslant 3$, $|Y| \geqslant 1$ and $|Z| \geqslant 3$. Put $j = |X|$ and $m = |Y|$. Assume that

$$X = \{x_0, \ldots, x_{j-1}\} \quad \text{and} \quad Z = \{z_0, \ldots, z_{m-1}\}.$$

Moreover, put $x_j = x_0$ and $z_m = z_0$. Let S be the ternary system defined as follows: $V(S) = X \cup Y \cup Z$ and the following statement holds for all $u, v, w \in V(S)$: $uvSw$ if and only if one of the conditions (9)–(12) holds:

(9) $$u, v \in X \cup Y \quad \text{and} \quad u \neq v = w;$$

(10) $$u, v \in Y \cup Z \quad \text{and} \quad u \neq v = w;$$

(11) there exists f, $0 \leqslant f \leqslant j-1$, such that $u = x_f$, $v = x_{f+1}$ and $w \in Z$;

(12) there exists h, $0 \leqslant h \leqslant m-1$, such that $u = z_h$, $v = z_{h+1}$ and $w \in X$.

Obviously, S is a simple signpost system and its underlying graph is connected. It is clear that $(x_0 \ldots x_{j-1}x_j, z)$ and $(z_0 \ldots z_{m-1}z_m, x)$ are confusing circuits in S for every $z \in Z$ and every $x \in X$.

Let S be a simple signpost system. Assume that the underlying graph of S is connected. It was proved in [4] that there exists no confusing circuit in S if S satisfies the following axiom (ALT):

(ALT) if $uvSx$ and $xySy$, then $uvSy$ or $xySu$ or $yxSv$ for all $u, v, x, y \in V(S)$.

The assumption that the underlying graph of a considered signpost system is connected is not needed in the next proposition:

Proposition 4. *Let S be a smooth signpost system. Then there exists no confusing circuit in S.*

P r o o f. Suppose, to the contrary, that there exist $u_0, u_1, \ldots, u_k, v \in V(S)$ such that $k \geqslant 1$ and (8) is a confusing circuit in S. Then $u_0u_1 \ldots u_kSv$ and $u_0 = u_k$. By Lemma 4, $k \geqslant 3$. Since u_0u_1Sv, Proposition 3 implies that $u_0u_1Su_1$. Since $u_0 = u_k$, Proposition 3 implies that $\neg(u_0u_1Su_k)$. Then there exists j, $1 \leqslant j \leqslant k-1$, such that $u_0u_1Su_j$ and $\neg(u_0u_1Su_{j+1})$. Recall that u_0u_1Sv. Since S is smooth, we get $\neg(u_ju_{j+1}Sv)$, which contradicts the fact that $u_0u_1 \ldots u_kSv$. Thus the proposition is proved. □

Corollary 1. *Let S be a smooth signpost system, let $u_0, \ldots, u_i, v \in V(S)$, where $i \geqslant 1$, and let $u_0 \ldots u_i S v$. Then $(u_0, \ldots, u_i)$ is a path in the underlying graph of S.*

P r o o f. Let G denote the underlying graph of S. Clearly, $(u_0, \ldots, u_i)$ is a walk in G. Suppose, to the contrary, that $(u_0, \ldots, u_i)$ is not a path. Then there exist j and k such that $0 \leqslant j < k \leqslant i$ and $u_j = u_k$. Then $(u_j \ldots u_k, v)$ is a confusing circuit in S, which is a contradiction. □

Let S be a smooth signpost system. By a *path* in S we mean a sequence $(u_0, \ldots, u_k)$, where $k \geqslant 1$, $u_0, \ldots, u_k \in V(S)$ and $u_0 \ldots u_k S u_k$. As follows from Corollary 1, every path in S is a nontrivial path in the underlying graph of S. Let $v_0, \ldots, v_m, u, v, w \in V(S)$, where $m \geqslant 1$; if $(v_0, \ldots, v_m)$ is a path in S, $u = v_0$, $v = v_1$ and $w = v_m$, then we say that $(v_0, \ldots, v_m)$ is an $uv - w$ path in S.

Theorem 1. *Let S be a smooth signpost system, and let $u, v, w \in V(S)$. If $uvSw$, then there exists an $uv - w$ path in S.*

P r o o f. Suppose, to the contrary, that there exist no $uv - w$ path in S. Then $v \neq w$. As follows from (SIG3), there exists an infinite sequence

$$v_0, v_1, v_2, \ldots$$

of elements of $V(S)$ such that $v_0 = u$, $v_1 = v$ and

$$v_0 \ldots v_i S w \text{ and } v_i \neq w \text{ for each } i = 1, 2, 3, \ldots.$$

Since $V(S)$ is finite, there exist distinct j and k such that $0 \leqslant j < k$ and $v_j = v_k$. Since $v_j \ldots v_k S w$, we see that $(v_j \ldots v_k, w)$ is a confusing circuit in S, which contradicts Proposition 4. Thus the theorem is proved. □

Remark. Theorem 1 can be interpreted as follows: Let S be a signpost system, let G denote the underlying graph of S, let $u, v, w \in V(S)$ and $uvSw$. If we know that S is smooth, then we are certain that the signpost (u, v, w) shows a path going from u to w in G; the direction (in u) of this path is determined by v.

The next lemma was proved in [4]:

Lemma 5 (see Lemma 2 in [4])**.** *Let S be a simple signpost system, and let $u_0, \ldots, u_{i-1}, u_i \in V(S)$, where $i \geqslant 1$. If $u_0 \ldots u_{i-1} u_i S u_i$, then $u_i u_{i-1} \ldots u_0 S u_0$.*

Hint for the proof. We proceed by induction on i. If $i = 1$, then we use (6). If $i \geqslant 2$, then we combine the induction hypothesis with Lemma 2 and (6). □

Corollary 2. *Let S be a simple smooth signpost system, and let $u_0, \dots, u_{i-1}, u_i \in V(S)$, where $i \geqslant 1$. If $(u_0, \dots, u_{i-1}, u_i)$ is a path in S, then $(u_i, u_{i-1}, \dots, u_0)$ is a path in S.*

P r o o f. Proof is obvious. □

3. Extensions of simple smooth signpost systems

The next lemma will be used in the proof of the main result of this section.

Lemma 6. *Let S be a simple smooth signpost system, and let $u_0, \dots, u_i$, $v_0, \dots, v_j, x, y \in V(S)$, where $i, j \geqslant 1$. Assume that $u_0 = v_0$ and $u_i = v_j$. If*

$$u_0 \dots u_i Sx \text{ and } v_0 \dots v_j Sy, \tag{13}$$

then

$$u_0 \dots u_i Sy \text{ and } v_0 \dots v_j Sx. \tag{14}$$

P r o o f. Let (13) hold. Since S is a smooth signpost system, Theorem 1 implies that there exist an $u_{i-1}u_i - x$ path in S and a $v_{j-1}v_j - y$ path in S.

If $u_i = x$, we put $m = i$. Let $u_i \neq x$. Then there exist $u_{i+1}, \dots u_m \in V(S)$, where $m \geqslant i + 1$, such that $u_m = x$ and $(u_{i-1}, u_i, \dots, u_m)$ is a path in S.

If $v_j = y$, we put $n = j$. Let $v_j \neq y$. Then there exist $v_{j+1}, \dots, v_n \in V(S)$, where $n \geqslant j + 1$, such that $v_n = y$ and $(v_{j-1}, v_j, \dots, v_n)$ is a path in S.

Clearly,

$$(u_0, \dots, u_{i-1}, u_i, \dots, u_m) \quad \text{and} \quad (v_0, \dots, v_{j-1}, v_j, \dots, v_n)$$

are paths in S. Corollary 2 implies that

$$(u_m, \dots, u_i, u_{i-1}, \dots, u_0) \quad \text{and} \quad (v_n, \dots, v_j, v_{j-1}, \dots, v_0)$$

are paths in S. Since $u_0 = v_0$ and $u_i = v_j$, we see that

$$(v_n, \dots, v_j = u_i, u_{i-1}, \dots, u_0) \quad \text{and} \quad (u_m, \dots, u_i = v_j, v_{j-1}, \dots v_0)$$

are paths in S and therefore, by Corollary 2,

$$(u_0, \dots, u_{i-1}, u_i = v_j, \dots, v_n) \quad \text{and} \quad (v_0, \dots, v_{j-1}, v_j = u_i, \dots, u_m)$$

are paths in S as well. Since $v_n = y$ and $u_m = x$, (14) follows, which completes the proof. □

Corollary 3. *Let S be a simple smooth signpost system, and let $u_0, \ldots, u_i, x \in V(S)$, where $i \geqslant 1$. If $u_0 \ldots u_i S x$ and $u_0 u_i S u_i$, then $u_0 u_i S x$.*

P r o o f. Proof is obvious. □

A graph G is called a *factor* of a graph H if $V(H) = V(G)$ and $E(G) \subseteq E(H)$.

Let S be a simple smooth signpost system, and let G denote the underlying graph of S. Consider a graph H such that G is a factor of H. By the *extension* of S to H we mean a ternary system S' with the following properties:

(a) $V(S') = V(S)$;

(b) if $u, v, w \in V(S')$, then $uvS'w$ if and only if u and v are adjacent vertices of H and there exist $u_0, \ldots, u_i \in V(S')$, where $i \geqslant 1$, such that $u_0 = u$, $u_i = v$ and $u_0 \ldots u_i S w$.

Clearly, the extension of S to H is defined uniquely.

As follows from Corollary 3, the extension of a simple smooth signpost system S to the underlying graph of S is identical to S.

Theorem 2. *Let S be a simple smooth signpost system, and let G denote the underlying graph of S. Consider a graph H such that G is a factor of H. Then the extension of S to H is a simple smooth signpost system and H is its underlying graph.*

P r o o f. Put $W = V(S)$. Let S' denote the extension of S to H. Obviously, S' is a ternary system with $V(S') = W$. The proof of the theorem will be divided into four parts. We will prove that

(1) S' is a signpost system,

(2) H is the underlying graph of S',

(3) S' is smooth, and

(4) S' is simple.

Part 1. Consider arbitrary $u, v, w \in W$ such that $uvS'w$. Then there exist $u_0, \ldots u_i \in W$, where $i \geqslant 1$, such that $u_0 = u$, $u_i = v$ and $u_0 \ldots u_i S w$. Hence $u_{i-1} u_i S w$. Recall that S is a smooth signpost system. If $u_i = w$, we put $m = i$. If $u_i \neq w$, then, by Theorem 1, there exist $u_{i+1}, \ldots u_m \in W$, where $m > i$, such that $u_m = w$ and $(u_{i-1}, u_i, \ldots, u_m)$ is a path in S. Hence $(u_0, \ldots, u_i, \ldots, u_m)$ is a path in S.

By Corollary 2, $(u_m, \ldots, u_i, \ldots, u_0)$ is a path in S as well. Hence $u_i \ldots u_0 S u_0$. This means that $vuS'u$. We see that S' satisfies (SIG1).

Assume that $vuS'w$. Then there exist $v_0, \ldots, v_j \in W$, where $j \geqslant 1$, such that $v_0 = v$, $v_j = u$ and $v_0 \ldots v_j S w$. Since $u_i = v_0$ and $v_j = u_0$, we see that $(u_0 \ldots u_i v_1 \ldots v_j, w)$ is a confusing circuit in S, which contradicts Proposition 4. Hence $\neg(vuS'w)$. We see that S' satisfies (SIG2).

Consider arbitrary $u^*, v^* \in W$ such that $u^* \neq v^*$. Then there exists $r \in W$ such that $u^* r S v^*$. Since $u*$ and r are adjacent vertices of H, we get $u^* r S' v^*$. We see that S' satisfies (SIG3). Therefore, S' is a signpost system.

Part 2. Obviously, $V(H) = W$. Consider arbitrary $u, v \in W$, $u \neq v$. As follows from the definition of S', if $uvS'v$, then $uv \in E(H)$. Conversely, let $uv \in E(H)$. As follows from (SIG3), there exists $r \in W$ such that $urS'v$. By Theorem 1, there exists an $ur - v$ path in S. This implies that there exist $u_0, \ldots, u_k$, where $k \geqslant 1$, such that $u_0 = u$, $u_1 = r$, $u_k = v$ and $u_0 \ldots u_k S u_k$. Thus $u_0 u_k S' u_k$; we have $uvS'v$. Therefore, H is the underlying graph of S'.

Part 3. Consider arbitrary $u, v, x, y \in W$ such that $uvS'x$, $uvS'z$ and $xyS'z$. Then $uv, xy \in E(H)$ and there exist $u_0, \ldots, u_i, v_0, \ldots, v_j, x_0, \ldots, x_k \in W$, where $i, j, k \geqslant 1$, such that $u_0 = u$, $u_i = v$, $v_0 = u, \ldots v_j = v$, $x_0 = x$, $x_k = y$,

$$u_0 \ldots u_i S x, v_0 \ldots v_j S z \quad \text{and} \quad x_0 \ldots x_k S z.$$

Since S is a simple smooth signpost system, Lemma 6 implies that $u_0 \ldots u_i S z$. By Proposition 1, $u_0 \ldots u_i S x_k$. Recall that $uv \in E(H)$. Hence $uvS'y$. We see that S' satisfies (SMO). Therefore, S' is smooth.

Part 4. Consider arbitrary $u, v, x, y \in W$ such that $uvS'x$ and $xyS'v$. Then $uv, xy \in E(H)$ and there exist $u_0, \ldots u_i, x_0, \ldots, x_j \in W$, where $i, j \geqslant 1$, such that $u_0 = v$, $u_i = u$, $x_0 = x$, $x_j = y$, $u_i \ldots u_0 S x_j$ and $x_0 \ldots x_j S u_0$. Since S is simple, Proposition 2 implies that $u_i \ldots u_0 S x_j$ and $x_0 \ldots x_j S u_i$. Recall that $uv, xy \in E(H)$. Hence $uvS'y$ and $xyS'u$. We see that S' satisfies (SIM1) and (SIM2). Therefore, S' is simple. □

4. Connected graphs

Let G be a connected graph. By the *step system* of G we mean the ternary system S such that $V(S) = V(G)$ and

$$uvSw \text{ if and only if } d(u,v) = 1 \text{ and } d(v,w) = d(u,w) - 1 \text{ for all } u, v, w \in V(G),$$

where $d(x, y)$ denotes the distance between vertices x and y in G. It is easy to show that the step system of a connected graph is a simple signpost system.

Lemma 7. *If G is a tree, then the step system of G is smooth.*

Proof. Proof is simple. □

As was proved in [3] (and also in [5]), a simple signpost system is the step system of a connected graph if and only if

(15) the underlying graph of S is connected

and S satisfies the axiom (ALT) and a certain pair of additional axioms. Almost the same result was proved in [6]; but the proof in [6] is very different both from the proof in [3] and the proof in [5]; the pair of additional axioms was replaced by the following axiom (COM) in [6]:

(COM) if $uvSx$, $vuSy$ and $xySu$, then $yxSv$ for all $u, v, x, y \in V(S)$.

The axioms (SIG1), (SIG2), (SIG3), (SIM1), (SIM2), (ALT) and (COM) (and, of course, the axiom (SWO)) could be formulated in a language of the first-order logic. It is unknown whether the condition (15) can be replaced by a finite set of axioms of the same kind.

The step systems of connected graphs of two special classes were characterized without help of the condition (15) in [2]; one of these classes is the class of median graphs (for the notion of a median graph the reader is referred to [1]); note that every tree is a median graph.

As was shown in [6], the step system of every median graph is smooth. On the other hand, if G is a connected graph that contains an induced K(2, 3), then the step system of G is not smooth.

The following theorem is the main result of the present paper.

Theorem 3. *Let G be a graph. Then the following statements are equivalent:*
(a) *G is connected;*
(b) *G is the underlying graph of a simple smooth signpost system;*
(c) *G is the underlying graph of a smooth signpost system.*

P r o o f. (a) → (b): Let G be connected. Consider an arbitrary spanning tree T of G. Let S_T denote the step system of T. Then S_T is a simple signpost system. By Lemma 7, S_T is smooth. Let S denote the extension of S_T to G. By Theorem 2, S is a simple smooth signpost system and the underlying graph of S is G.

(b) → (c): Obvious.

(c) → (a): Let G be the underlying graph of a smooth signpost system S. Consider arbitrary $u, v \in V(S)$ such that $u \neq v$. By (SIG3), there exists $t \in V(S)$ such that $utSv$. Theorem 1 implies that there exists an $ut - v$ path in S. Clearly, there exists a path connecting u and v in G. This implies that G is connected. □

References

[1] *H. M. Mulder*: The Interval Function of a Graph. Math. Centre Tracts 132. Math. Centre, Amsterdam, 1980.

[2] *H. M. Mulder and L. Nebeský*: Modular and median signpost systems and their underlying graphs. Discussiones Mathematicae Graph Theory *23* (2003), 309–324.

[3] *L. Nebeský*: Geodesics and steps in a connected graph. Czechoslovak Math. J. *47(122)* (1997), 149–161.

[4] *L. Nebeský*: An axiomatic approach to metric properties of connected graphs. Czechoslovak Math. J. *50(125)* (2000), 3–14.

[5] *L. Nebeský*: A theorem for an axiomatic aproach to metric properties of graphs. Czechoslovak Math. J. *50(125)* (2000), 121–133.

[6] *L. Nebeský*: On properties of a graph that depend on its distance function. Czechoslovak Math. J. *54(129)* (2004), 445–456.

Author's address: Univerzita Karlova v Praze, Filozofická fakulta, nám. J. Palacha 2, 116 38 Praha 1, Czech Republic, e-mail: `Ladislav.Nebesky@ff.cuni.cz`.