Czechoslovak Mathematical Journal, 69 (144) (2019), 353–363

A DIOPHANTINE INEQUALITY WITH FOUR SQUARES AND ONE kTH POWER OF PRIMES

Quanwu Mu, Minhui Zhu, Ping Li, Xi'an

Received July 3, 2017. Published online November 19, 2018.

Abstract. Let $k \geqslant 5$ be an odd integer and η be any given real number. We prove that if λ_1, λ_2, λ_3, λ_4, μ are nonzero real numbers, not all of the same sign, and λ_1/λ_2 is irrational, then for any real number σ with $0 < \sigma < 1/(8\vartheta(k))$, the inequality

$$|\lambda_1 p_1^2 + \lambda_2 p_2^2 + \lambda_3 p_3^2 + \lambda_4 p_4^2 + \mu p_5^k + \eta| < \Big(\max_{1 \leqslant j \leqslant 5} p_j\Big)^{-\sigma}$$

has infinitely many solutions in prime variables $p_1, p_2, \ldots, p_5$, where $\vartheta(k) = 3 \times 2^{(k-5)/2}$ for $k = 5, 7, 9$ and $\vartheta(k) = [(k^2 + 2k + 5)/8]$ for odd integer k with $k \geqslant 11$. This improves a recent result in W. Ge, T. Wang (2018).

Keywords: Diophantine inequalities; Davenport-Heilbronn method; prime

MSC 2010: 11D75, 11P55

1. Introduction

In 1937, Vinogradov [23] proved that every sufficiently large odd integer is a sum of three primes. Later, Hua [11] refined Vinogradov's result and showed that all sufficiently large odd integers are sums of two primes and a kth power of a prime, where k is any given positive integer. In [11], Hua also proved that all sufficiently large odd integers satisfying some necessary congruence conditions can be represented

This work was supported by the National Natural Science Foundation of China (No. 11771333), the Natural Science Basic Research Plan in Shaanxi Province of China (No. 2016JM1013), the Scientific Research Program Funded by Shaanxi Provincial Education Department (No. 17JK0341) and the Quality Improvement Project for Basic Curriculum (Functions of Complex Variables and Integral Transformation) of Xi'an Polytechnic University.

DOI: 10.21136/CMJ.2018.0316-17

in the form of four squares of primes and a kth power of a prime. It is of some interest to consider the analogous form for Diophantine inequalities. Some authors obtained many significant results in this direction, see [1], [2], [6], [8], [9], [13], [14], [15], [16], [19], [20], [21] for details. In [14], Li and Wang established the following theorem.

Theorem 1.1. *Let $k \geqslant 3$ be a fixed integer and η be any given real number. Suppose that λ_1, λ_2, λ_3, λ_4, μ are nonzero real numbers, not all of the same sign, and λ_1/λ_2 is irrational. Then the inequality*

$$|\lambda_1 p_1^2 + \lambda_2 p_2^2 + \lambda_3 p_3^2 + \lambda_4 p_4^2 + \mu p_5^k + \eta| < \Big(\max_{1\leqslant j\leqslant 5} p_j\Big)^{-\sigma} \tag{1.1}$$

has infinitely many solutions in prime variables $p_1, p_2, \ldots, p_5$ for $0 < \sigma < 1/(3k2^k)$.

In [17], we improved the above result and showed that under the same assumptions as in Theorem 1.1, inequality (1.1) has infinitely many solutions in prime variables $p_1, p_2, \ldots, p_5$, where $0 < \sigma < 1/16$ for $k = 3$, $0 < \sigma < 5/(3k2^k)$ for $4 \leqslant k \leqslant 5$, and $0 < \sigma < 40/(21k2^k)$ for $k \geqslant 6$. The proof is based on the method of Languasco and Zaccagnini in [12], together with Heath-Brown's improvement on Hua's lemma (see [4], Lemma 5 and [10], Theorem 2). Let

$$s(k) = \Big[\frac{k+1}{2}\Big], \quad \sigma(k) = \min(2^{s(k)-1}, \tfrac{1}{2}s(k)(s(k)+1)),$$

where $[x]$ denotes the largest integer not exceeding the real number x. Very recently, Ge and Wang [6] refined the result in [17]. They proved that under the same assumptions as in Theorem 1.1, inequality (1.1) has infinitely many solutions in prime variables $p_1, p_2, \ldots, p_5$ for $0 < \sigma < 1/(8\sigma(k))$ (see [6], Theorem 1.3).

The aim of the present paper is to further enlarge the range $0 < \sigma < 1/(8\sigma(k))$ for odd integer k with $k \geqslant 5$. The following theorem is proved.

Theorem 1.2. *Let $k \geqslant 5$ be an odd integer. Suppose that λ_1, λ_2, λ_3, λ_4, μ and η satisfy the same conditions as in Theorem 1.1. Then for any real number σ with $0 < \sigma < 1/(8\vartheta(k))$, inequality (1.1) has infinitely many solutions in prime variables $p_1, p_2, \ldots, p_5$, where*

$$\vartheta(k) = \begin{cases} 3 \times 2^{(k-5)/2} & \text{if } k = 5, 7, 9, \\ [(k^2+2k+5)/8] & \text{if } k \geqslant 11 \text{ and } 2 \nmid k. \end{cases} \tag{1.2}$$

With the help of Corollary 3.2 below, we obtain a wider major arc, this with the very recent work of Bourgain (see [3], Theorem 10) yields the desired conclusion.

2. Notation and preliminaries

The proof of Theorem 1.2 is dependent on the Davenport-Heilbronn circle method (see [22], Chapter 11). For each integer $j \geqslant 2$ set

$$\psi(j) = \begin{cases} 2^j & \text{when } 2 \leqslant j \leqslant 4, \\ j(j+1) & \text{when } j \geqslant 5. \end{cases} \tag{2.1}$$

In what follows, we use ε and δ to denote fixed positive constants which are arbitrarily small. The letter p, with or without subscript, always stands for a prime number. The letter k, except as specially provided, usually denotes an odd integer not less than 5. Since λ_1/λ_2 is irrational, we let q be a large enough denominator of a convergent to λ_1/λ_2. Put

$$X = q^2, \quad \mathcal{N}(X) = \sum_{\substack{\delta X \leqslant p_j^2 \leqslant X,\ 1 \leqslant j \leqslant 4,\ \delta X \leqslant p_5^k \leqslant X \\ |\lambda_1 p_1^2 + \lambda_2 p_2^2 + \lambda_3 p_3^2 + \lambda_4 p_4^2 + \mu p_5^k + \eta| < \tau}} 1,$$

$$\tau = X^{-1/(16\vartheta(k)) + 30\varepsilon}, \quad K_\tau(\alpha) = \begin{cases} \Big(\dfrac{\sin(\pi\tau\alpha)}{\pi\alpha}\Big)^2 & \text{when } \alpha \neq 0, \\ \tau^2 & \text{when } \alpha = 0, \end{cases}$$

$$S_j(\alpha) = \sum_{\delta X \leqslant p^j \leqslant X} (\log p) e(\alpha p^j),$$

$$I(\tau, \eta, \mathfrak{X}) = \int_{\mathfrak{X}} \prod_{j=1}^{4} S_2(\lambda_j \alpha)\, S_k(\mu\alpha) e(\alpha\eta) K_\tau(\alpha)\, \mathrm{d}\alpha,$$

where $e(\alpha) = \mathrm{e}^{2\pi \mathrm{i}\alpha}$, $\mathfrak{X}$ denotes any measurable subset of $\mathbb{R}$ and $\vartheta(k)$ is defined by (1.2). For the Dirichlet kernel $K_\tau(\alpha)$ we have the trivial estimate

$$K_\tau(\alpha) \ll \min(\tau^2,\ |\alpha|^{-2}). \tag{2.2}$$

It follows from Lemma 4 of Davenport and Heilbronn [5] that

$$\int_{-\infty}^{\infty} e(xy) K_\tau(x)\, \mathrm{d}x = \max(0, \tau - |y|). \tag{2.3}$$

Thus,

$$
\begin{aligned}
(2.4)\qquad \mathcal{N}(X) &\geqslant \frac{1}{\tau} \sum_{\substack{\delta X \leqslant p_j^2 \leqslant X \\ 1 \leqslant j \leqslant 4 \\ \delta X \leqslant p_5^k \leqslant X}} \max(0, \tau - |\lambda_1 p_1^2 + \lambda_2 p_2^2 + \lambda_3 p_3^2 + \lambda_4 p_4^2 + \mu p_5^k + \eta|) \\
&\geqslant \frac{1}{\tau (\log X)^5} \sum_{\substack{\delta X \leqslant p_j^2 \leqslant X \\ 1 \leqslant j \leqslant 4 \\ \delta X \leqslant p_5^k \leqslant X}} \prod_{j=1}^{5} \log p_j \\
&\quad \times \max(0, \tau - |\lambda_1 p_1^2 + \lambda_2 p_2^2 + \lambda_3 p_3^2 + \lambda_4 p_4^2 + \mu p_5^k + \eta|) \\
&= \frac{1}{\tau (\log X)^5} \sum_{\substack{\delta X \leqslant p_j^2 \leqslant X \\ 1 \leqslant j \leqslant 4 \\ \delta X \leqslant p_5^k \leqslant X}} \prod_{j=1}^{5} \log p_j \\
&\quad \times \int_{-\infty}^{\infty} e((\lambda_1 p_1^2 + \lambda_2 p_2^2 + \lambda_3 p_3^2 + \lambda_4 p_4^2 + \mu p_5^k + \eta)\alpha) K_\tau(\alpha) \, \mathrm{d}\alpha \\
&= \frac{1}{\tau (\log X)^5} \, I(\tau, \eta, \mathbb{R}).
\end{aligned}
$$

To prove Theorem 1.2, it suffices to establish the estimate $I(\tau, \eta, \mathbb{R}) \gg \tau^2 X^{1+1/k}$. For this purpose, we split the real line into three parts

$$
\mathfrak{M} = \{\alpha \colon |\alpha| \leqslant \varphi\}, \quad \mathfrak{m} = \{\alpha \colon \varphi < |\alpha| \leqslant \xi\}, \quad \mathfrak{t} = \{\alpha \colon |\alpha| > \xi\},
$$

where $\varphi = X^{-1/(2k)-\varepsilon}$, $\xi = \tau^{-2} X^{3\varepsilon}$. Usually, these sets are called the major arc, the minor arcs and the trivial arcs, respectively. Therefore

$$
(2.5)\qquad I(\tau, \eta, \mathbb{R}) = I(\tau, \eta, \mathfrak{M}) + I(\tau, \eta, \mathfrak{m}) + I(\tau, \eta, \mathfrak{t}).
$$

It should be noted that the major arc $\mathfrak{M}$ is wider than that of [6]. In what follows, we shall show that

$$
|I(\tau, \eta, \mathfrak{M})| \gg \tau^2 X^{1+1/k}, \quad |I(\tau, \eta, \mathfrak{m})| \ll \tau^2 X^{1+1/k-\varepsilon}, \quad |I(\tau, \eta, \mathfrak{t})| \ll \tau^2 X^{1+1/k-\varepsilon}.
$$

3. The major arc

Let $\mathbf{M} = \{\alpha\colon |\alpha| \leqslant X^{-1+5/(6k)-\varepsilon}\}$, then $\mathbf{M} \subset \mathfrak{M}$. In [17], Section 3, we have proved that

$$|I(\tau,\eta,\mathbf{M})| \gg \tau^2 X^{1+1/k}. \tag{3.1}$$

The conditions 'λ_1, λ_2, λ_3, λ_4, μ are nonzero real numbers, not all of the same sign' play an important role in the proof of (3.1), see [17], pages 485–486 for details. It remains to discuss the estimate for $|I(\tau,\eta,\mathfrak{M}\setminus\mathbf{M})|$.

Lemma 3.1 (see [7], Theorem 1)**.** *Let j be an integer with $j \geqslant 2$, and $N \geqslant 2$. Suppose that a and q are integers with*

$$|q\alpha - a| \leqslant \frac{1}{q}, \quad (a,\ q) = 1, \quad q \geqslant 1. \tag{3.2}$$

Then for any $\varepsilon > 0$,

$$\sum_{p\leqslant N} (\log p) e(\alpha p^j) \ll N^{1+\varepsilon}\Big(\frac{1}{q} + \frac{1}{N^{1/2}} + \frac{q}{N^j}\Big)^{4^{1-j}}. \tag{3.3}$$

Corollary 3.2. *Suppose that $X^{-1+5/(6k)-\varepsilon} \leqslant |\alpha| \leqslant X^{-1/(2k)-\varepsilon}$. Then for any given nonzero real μ and $\varepsilon > 0$ we have*

$$|S_k(\mu\alpha)| \ll X^{1/k(1-1/2\times 4^{1-k})+\varepsilon}. \tag{3.4}$$

The implicit constant in the $\ll$ notation depends on k, μ, δ.

Proof. Notice that

$$|S_k(\mu\alpha)| \leqslant \Big|\sum_{p\leqslant X^{1/k}} (\log p) e(\mu\alpha p^k)\Big| + \Big|\sum_{p\leqslant (\delta X)^{1/k}} (\log p) e(\mu\alpha p^k)\Big|. \tag{3.5}$$

Similarly to [9], Corollary 2, we take $\mu\alpha$ in place of α in (3.2), and take $q = [1/|\mu\alpha|]$, $a = \pm 1$ (the sign of a is the same as that for $\mu\alpha$), then (3.4) follows from (3.5) and (3.3). $\square$

By Corollary 3.2 and the arithmetic-geometric mean inequality, we get

$$
\begin{aligned}
(3.6)\qquad & |I(\tau,\eta,\mathfrak{M}\setminus\mathbf{M})| \\
&\leqslant \int_{\mathfrak{M}\setminus\mathbf{M}} |S_2(\lambda_1\alpha)S_2(\lambda_2\alpha)S_2(\lambda_3\alpha)S_2(\lambda_4\alpha)S_k(\mu\alpha)|K_\tau(\alpha)\,\mathrm{d}\alpha \\
&\ll \tau^2 \max_{\alpha\in\mathfrak{M}\setminus\mathbf{M}} |S_k(\mu\alpha)| \sum_{j=1}^{4} \int_{\mathfrak{M}\setminus\mathbf{M}} \bigl|S_2(\lambda_j\alpha)\bigr|^4\,\mathrm{d}\alpha \\
&\ll \tau^2 X^{1/k(1-1/2\times 4^{1-k})+\varepsilon} \int_0^1 \bigl|S_2(\alpha)\bigr|^4\,\mathrm{d}\alpha \\
&\ll \tau^2 X^{1+1/k-\varepsilon},
\end{aligned}
$$

where (2.2) and Hua's lemma (see [4], page 85) are used. Noting that $I(\tau,\eta,\mathfrak{M}) = I(\tau,\eta,\mathbf{M}) + I(\tau,\eta,\mathfrak{M}\setminus\mathbf{M})$, this with (3.1) and (3.6) implies

$$
(3.7)\qquad |I(\tau,\eta,\mathfrak{M})| \gg \tau^2 X^{1+1/k}.
$$

4. The minor arcs

Let $\widetilde{\mathfrak{m}} = \mathfrak{m}_1 \cup \mathfrak{m}_2$, where

$$
\mathfrak{m}_j = \{\alpha\in\mathfrak{m}\colon\ |S_2(\lambda_j\alpha)| \leqslant X^{7/16+2\varepsilon}\} \quad \text{for } j=1,2.
$$

To estimate the integral $I(\tau,\eta,\mathfrak{m})$, we need several lemmas.

Lemma 4.1. *Let j and s be positive integers with $s\leqslant j$. Then*

$$
(4.1)\qquad \int_0^1 |S_j(\alpha)|^{s(s+1)}\,\mathrm{d}\alpha \ll X^{s^2/j+\varepsilon}
$$

holds for all $\varepsilon>0$.

Proof. It follows from [3], Theorem 10 that

$$
(4.2)\qquad \int_0^1 \Bigl|\sum_{\delta X\leqslant x^j\leqslant X} e(\alpha x^j)\Bigr|^{s(s+1)}\,\mathrm{d}\alpha \ll X^{s^2/j+\varepsilon}.
$$

By considering the number of solutions of the underlying Diophantine equation and using (4.2), we obtain (4.1). $\square$

Lemma 4.2. *Let $j \geqslant 2$ be an integer. Suppose that λ and μ are nonzero real constants and k is an odd integer with $k \geqslant 5$. Then for any $\varepsilon > 0$ we have*

$$\int_{\mathbb{R}} |S_j(\lambda\alpha)|^{\psi(j)} K_\tau(\alpha)\,\mathrm{d}\alpha \ll \tau X^{\psi(j)/j-1+\varepsilon}, \tag{4.3}$$

$$\int_{\mathbb{R}} |S_2(\lambda\alpha)|^2 |S_k(\mu\alpha)|^{2\vartheta(k)} K_\tau(\alpha)\,\mathrm{d}\alpha \ll \tau X^{2\vartheta(k)/k+\varepsilon}, \tag{4.4}$$

where $\psi(j)$ and $\vartheta(k)$ are defined by (2.1) *and* (1.2)*, respectively. The implicit constant in the $\ll$ notation of* (4.3) *depends on λ, j, and the implicit constant in the $\ll$ notation of* (4.4) *depends on k, λ, μ.*

P r o o f. For (4.3), see [18], Lemma 4.5 for details. It remains to prove (4.4). Let $a = (k-1)/2$, $b = (k+1)/2$.

We first consider the case of $k \geqslant 11$, $2 \nmid k$, recalling that $\vartheta(k) = [(k^2+2k+5)/8]$ in this case. When $k \equiv 1 \pmod 4$, we have

$$\vartheta(k) = \frac{k^2+2k+5}{8} = \frac{a(a+1)+b(b+1)}{4} + \frac{1}{2}.$$

It follows from the Cauchy-Schwarz inequality and Lemma 4.1 that

$$\begin{aligned}\int_0^1 |S_k(\alpha)|^{2\vartheta(k)}\,\mathrm{d}\alpha &\ll X^{1/k}\int_0^1 |S_k(\alpha)|^{(a(a+1)+b(b+1))/2}\,\mathrm{d}\alpha \\ &\ll X^{1/k}\biggl(\int_0^1 |S_k(\alpha)|^{a(a+1)}\biggr)^{1/2}\biggl(\int_0^1 |S_k(\alpha)|^{b(b+1)}\biggr)^{1/2} \\ &\ll X^{1/k}(X^{a^2/k+\varepsilon})^{1/2}(X^{b^2/k+\varepsilon})^{1/2} \\ &\ll X^{(k^2+5)/(4k)+\varepsilon} \ll X^{2\vartheta(k)/k-1/2+\varepsilon},\end{aligned} \tag{4.5}$$

where the trivial upper bound $S_k(\alpha) \ll X^{1/k}$ is used. When $k \equiv 3 \pmod 4$, we have

$$\vartheta(k) = \frac{(k+1)^2}{8} = \frac{a(a+1)+b(b+1)}{4}.$$

By a similar argument as that in (4.5), we also obtain

$$\int_0^1 |S_k(\alpha)|^{2\vartheta(k)}\,\mathrm{d}\alpha \ll X^{2\vartheta(k)/k-1/2+\varepsilon}. \tag{4.6}$$

It follows from (2.3) that

$$\int_{\mathbb{R}} |S_2(\lambda\alpha)|^2 |S_k(\mu\alpha)|^{2\vartheta(k)} K_\tau(\alpha)\,\mathrm{d}\alpha \ll \tau\Sigma, \tag{4.7}$$

where Σ denotes the number of solutions of

$$|\mu(p_1^k+\ldots+p_{\vartheta(k)}^k-p_{\vartheta(k)+1}^k-\ldots-p_{2\vartheta(k)}^k)+\lambda(p_{2\vartheta(k)+1}^2-p_{2\vartheta(k)+2}^2)|<\tau$$

with $p_i^k\in[\delta X,X]$ for $1\leqslant i\leqslant 2\vartheta(k)$, and $p_j^2\in[\delta X,X]$ for $2\vartheta(k)+1\leqslant j\leqslant 2\vartheta(k)+2$. Note that $\tau\to 0$ as $X\to\infty$. When $p_{2\vartheta(k)+1}\neq p_{2\vartheta(k)+2}$, the values of $p_1,p_2,\ldots,p_{2\vartheta(k)}$ determine the values of $p_{2\vartheta(k)+1}$ and $p_{2\vartheta(k)+2}$ with at most X^ε possibilities; these solutions contribute $\ll X^{2\vartheta(k)/k+\varepsilon}$ to Σ. When $p_{2\vartheta(k)+1}=p_{2\vartheta(k)+2}$, we get

$$p_1^k+\ldots+p_{\vartheta(k)}^k-p_{\vartheta(k)+1}^k-\ldots-p_{2\vartheta(k)}^k=0. \tag{4.8}$$

By (4.5) and (4.6), it follows that equation(4.8) has $O(X^{2\vartheta(k)/k-1/2+\varepsilon})$ solutions in primes $p_1,p_2,\ldots,p_{2\vartheta(k)}$. In this case, these solutions also contribute $\ll X^{2\vartheta(k)/k+\varepsilon}$ to Σ. Thus, we get $\Sigma\ll X^{2\vartheta(k)/k+\varepsilon}$; this with (4.7) yields (4.4).

In the cases of $k=5,7,9$, noting that $\vartheta(k)=3\times 2^{(k-5)/2}=2^{a-2}+2^{b-2}$, we can also prove (4.6) using the Cauchy-Schwarz inequality and Hua's lemma. In a similar manner as above, we can prove (4.4). This completes the proof of Lemma 4.2. $\square$

From the arithmetic-geometric mean inequality, Hölder's inequality and Lemma 4.2, we get

$$\begin{aligned}
I(\tau,\eta,\mathfrak{m}_1)&\ll\sum_{j=2}^4\int_{\mathfrak{m}_1}|S_2(\lambda_1\alpha)||S_2(\lambda_j\alpha)|^3|S_k(\mu\alpha)|K_\tau(\alpha)\,\mathrm{d}\alpha\\
&\ll\Big(\sup_{\alpha\in\mathfrak{m}_1}|S_2(\lambda_1\alpha)|\Big)^{1/\vartheta(k)}\Big(\int_{\mathbb{R}}|S_2(\lambda_1\alpha)|^4K_\tau(\alpha)\,\mathrm{d}\alpha\Big)^{1/4-1/(2\vartheta(k))}\\
&\quad\times\Big(\int_{\mathbb{R}}|S_2(\lambda_1\alpha)|^2|S_k(\mu\alpha)|^{2\vartheta(k)}K_\tau(\alpha)\,\mathrm{d}\alpha\Big)^{1/(2\vartheta(k))}\\
&\quad\times\sum_{j=2}^4\Big(\int_{\mathbb{R}}|S_2(\lambda_j\alpha)|^4K_\tau(\alpha)\,\mathrm{d}\alpha\Big)^{3/4}\\
&\ll(X^{7/16+2\varepsilon})^{1/\vartheta(k)}(\tau X^{1+\varepsilon})^{1/4-1/(2\vartheta(k))}(\tau X^{2\vartheta(k)/k+\varepsilon})^{1/(2\vartheta(k))}(\tau X^{1+\varepsilon})^{3/4}\\
&\ll\tau X^{1+1/k-1/(16\vartheta(k))+4\varepsilon}\ll\tau^2X^{1+1/k-\varepsilon}.
\end{aligned}$$

By symmetry, the same bound holds for $\mathfrak{m}_2$ in place of $\mathfrak{m}_1$. This implies that

$$I(\tau,\eta,\widetilde{\mathfrak{m}})\ll\tau^2X^{1+1/k-\varepsilon}. \tag{4.9}$$

It therefore remains to discuss the set $\mathfrak{m}^*=\mathfrak{m}\setminus\widetilde{\mathfrak{m}}$, in which

$$|S_2(\lambda_1\alpha)|>X^{7/16+2\varepsilon},\quad |S_2(\lambda_2\alpha)|>X^{7/16+2\varepsilon},\quad X^{-1/(2k)-\varepsilon}<|\alpha|\leqslant\tau^{-2}X^{3\varepsilon}$$

hold simultaneously. By a familiar dyadic dissection argument, we divide $\mathfrak{m}^*$ into at most $\ll \log^3 X$ disjoint sets $E(Z_1, Z_2, y)$. For $\alpha \in E(Z_1, Z_2, y)$ we have

$$Z_1 < |S_2(\lambda_1\alpha)| \leqslant 2Z_1, \quad Z_2 < |S_2(\lambda_2\alpha)| \leqslant 2Z_2, \quad y < |\alpha| \leqslant 2y,$$

where $Z_1 = 2^{k_1}X^{7/16+2\varepsilon}$, $Z_2 = 2^{k_2}X^{7/16+2\varepsilon}$ and $y = 2^{k_3}X^{-1/(2k)-\varepsilon}$ for some non-negative integers k_1, k_2, k_3.

For simplicity, we take the notation $\mathscr{A}$ as a shortcut for $E(Z_1, Z_2, y)$, and let $m(\mathscr{A})$ denote the Lebesgue measure of $\mathscr{A}$.

Lemma 4.3. *We have*

$$m(\mathscr{A}) \ll yX^{5/2+8\varepsilon}(Z_1Z_2)^{-4}.$$

P r o o f. See [17], Lemma 6. □

By (2.2), the arithmetic-geometric mean inequality and Hölder's inequality, we have

$$\begin{aligned}
I(\tau,\eta,\mathscr{A}) &\ll \sum_{j=3}^{4}\int_{\mathscr{A}} |S_2(\lambda_1\alpha)S_2(\lambda_2\alpha)||S_2(\lambda_j\alpha)|^2|S_k(\mu\alpha)|K_\tau(\alpha)\,\mathrm{d}\alpha \\
&\ll \bigg(\int_{\mathscr{A}} |S_2(\lambda_1\alpha)S_2(\lambda_2\alpha)|^4K_\tau(\alpha)\,\mathrm{d}\alpha\bigg)^{1/4}\bigg(\int_{\mathbb{R}} |S_k(\mu\alpha)|^{\psi(k)}K_\tau(\alpha)\,\mathrm{d}\alpha\bigg)^{1/\psi(k)} \\
&\quad\times\bigg(\int_{\mathscr{A}} K_\tau(\alpha)\,\mathrm{d}\alpha\bigg)^{1/4-1/\psi(k)}\sum_{j=3}^{4}\bigg(\int_{\mathbb{R}} |S_2(\lambda_j\alpha)|^4K_\tau(\alpha)\,\mathrm{d}\alpha\bigg)^{1/2} \\
&\ll ((Z_1Z_2)^4m(\mathscr{A})\min(\tau^2,y^{-2}))^{1/4}(\tau X^{\psi(k)/k-1+\varepsilon})^{1/\psi(k)} \\
&\quad\times(\min(\tau^2,y^{-2})m(\mathscr{A}))^{1/4-1/\psi(k)}(\tau X^{1+\varepsilon})^{1/2} \\
&\ll \tau^{1/2+1/\psi(k)}(y\min(\tau^2,y^{-2}))^{1/2-1/\psi(k)}X^{7/8+1/k+3\varepsilon} \\
&\ll \tau X^{7/8+1/k+3\varepsilon} \ll \tau^2X^{1+1/k-2\varepsilon},
\end{aligned}$$

where Lemmas 4.2–4.3 and the bounds $Z_j \geqslant X^{7/16+2\varepsilon}$ $(j = 1, 2)$ are used. Thus,

$$I(\tau,\eta,\mathfrak{m}^*) \ll (\log^3 X)\max_{\mathscr{A}}|I(\tau,\eta,\mathscr{A})| \ll \tau^2X^{1+1/k-\varepsilon}. \tag{4.10}$$

It follows from (4.9) and (4.10) that

$$I(\tau,\eta,\mathfrak{m}) \ll \tau^2X^{1+1/k-\varepsilon}. \tag{4.11}$$

5. The trivial arcs

The proof of $|I(\tau,\eta,\mathfrak{t})| \ll \tau^2 X^{1+1/k-\varepsilon}$ is almost identical to that of inequality (25) in [17]. We list it for the sake of completeness.

$$
\begin{aligned}
\big|I(\tau,\eta,\mathfrak{t})\big| &\ll \int_{\xi}^{\infty} |S_2(\lambda_1\alpha)S_2(\lambda_2\alpha)S_2(\lambda_3\alpha)S_2(\lambda_4\alpha)S_k(\mu\alpha)\big|K_\tau(\alpha)\,\mathrm{d}\alpha \\
&\ll X^{1/k}\sum_{j=1}^{4}\int_{\xi}^{\infty} |S_2(\lambda_j\alpha)|^4 K_\tau(\alpha)\,\mathrm{d}\alpha \\
&\ll X^{1/k}\sum_{j=1}^{4}\int_{|\lambda_j|\xi}^{\infty} \frac{|S_2(\alpha)|^4}{\alpha^2}\,\mathrm{d}\alpha \\
&\ll X^{1/k}\sum_{j=1}^{4}\sum_{n\geqslant|\lambda_j|\xi} \frac{1}{(n-1)^2}\int_{n-1}^{n} |S_2(\alpha)|^4\,\mathrm{d}\alpha \\
&\ll \frac{X^{1/k}X^{1+\varepsilon}}{\xi} \ll \tau^2 X^{1+1/k-\varepsilon}.
\end{aligned}
\tag{5.1}
$$

6. Completion of the proof

By (3.7), (4.11), (5.1) and (2.5), we get $I(\tau,\eta,\mathbb{R}) \gg \tau^2 X^{1+1/k}$. It follows from (2.4) that

$$
\mathcal{N}(X) \gg \tau X^{1+1/k}(\log X)^{-5} \gg X^{1+1/k-1/(16\vartheta(k))+\varepsilon}.
$$

Recalling that λ_1/λ_2 is irrational, q is a large enough denominator of a convergent to λ_1/λ_2 and $X = q^2$. When $q \to \infty$, we have $X \to \infty$; this implies $\mathcal{N}(X) \to \infty$. The value of τ and $\max p_j \asymp X^{1/2}$ give the desired range of σ on the right-hand side of (1.1). This completes the proof of Theorem 1.2.

Acknowledgment. The authors are very grateful to the referee for many valuable suggestions.

References

[1] *A. Baker*: On some diophantine inequalities involving primes. J. Reine Angew. Math. *228* (1967), 166–181. zbl MR doi

[2] *R. C. Baker, G. Harman*: Diophantine approximation by prime numbers. J. Lond. Math. Soc., II. Ser. *25* (1982), 201–215. zbl MR doi

[3] *J. Bourgain*: On the Vinogradov mean value. Proc. Steklov Inst. Math. *296* (2017), 30–40; translated from Tr. Mat. Inst. Steklova *296* (2017), 36–46. zbl MR doi

[4] *R. J. Cook*: The value of additive forms at prime arguments. J. Théor. Nombres Bordx. *13* (2001), 77–91. zbl MR doi
[5] *H. Davenport, H. Heilbronn*: On indefinite quadratic forms in five variables. J. Lond. Math. Soc. *21* (1946), 185–193. zbl MR doi
[6] *W. Ge, T. Wang*: On Diophantine problems with mixed powers of primes. Acta Arith. *182* (2018), 183–199. zbl MR doi
[7] *G. Harman*: Trigonometric sums over primes I. Mathematika *28* (1981), 249–254. zbl MR doi
[8] *G. Harman*: Diophantine approximation by prime numbers. J. Lond. Math. Soc., II. Ser. *44* (1991), 218–226. zbl MR doi
[9] *G. Harman*: The values of ternary quadratic forms at prime arguments. Mathematika *51* (2004), 83–96. zbl MR doi
[10] *D. R. Heath-Brown*: Weyl's inequality, Hua's inequality, and Waring's problem. J. Lond. Math. Soc., II. Ser. *38* (1988), 216–230. zbl MR doi
[11] *L.-K. Hua*: Some results in additive prime-number theory. Q. J. Math., Oxf. Ser. *9* (1938), 68–80. zbl MR doi
[12] *A. Languasco, A. Zaccagnini*: A Diophantine problem with a prime and three squares of primes. J. Number Theory *132* (2012), 3016–3028. zbl MR doi
[13] *A. Languasco, A. Zaccagnini*: A Diophantine problem with prime variables. Highly Composite: Papers in Number Theory (V. Kumar Murty, R. Thangadurai, eds.). Ramanujan Mathematical Society Lecture Notes Series 23, Ramanujan Mathematical Society, Mysore, 2016, pp. 157–168. zbl MR
[14] *W. Li, T. Wang*: Diophantine approximation with four squares and one k-th power of primes. J. Math. Sci. Adv. Appl. *6* (2010), 1–16. zbl MR
[15] *W. Li, T. Wang*: Diophantine approximation with two primes and one square of prime. Chin. Q. J. Math. *27* (2012), 417–423. zbl
[16] *K. Matomäki*: Diophantine approximation by primes. Glasg. Math. J. *52* (2010), 87–106. zbl MR doi
[17] *Q. Mu*: Diophantine approximation with four squares and one kth power of primes. Ramanujan J. *39* (2016), 481–496. zbl MR doi
[18] *Q. Mu*: One Diophantine inequality with unlike powers of prime variables. Int. J. Number Theory *13* (2017), 1531–1545. zbl MR doi
[19] *Q. Mu, Y. Qu*: A Diophantine inequality with prime variables and mixed power. Acta Math. Sin., Chin. Ser. *58* (2015), 491–500. (In Chinese.) zbl MR
[20] *K. Ramachandra*: On the sums $\sum_{j=1}^{K} \lambda_j f_j(p_j)$. J. Reine Angew. Math. *262/263* (1973), 158–165. zbl MR doi
[21] *R. C. Vaughan*: Diophantine approximation by prime numbers. I. Proc. Lond. Math. Soc., III. Ser. *28* (1974), 373–384. zbl MR doi
[22] *R. C. Vaughan*: The Hardy-Littlewood Method. Cambridge Tracts in Mathematics 125, Cambridge University Press, Cambridge, 1997. zbl MR doi
[23] *I. M. Vinogradov*: Representation of an odd number as a sum of three primes. C. R. (Dokl.) Acad. Sci. URSS, n. Ser. *15* (1937), 169–172. zbl

Authors' address: Quanwu Mu (corresponding author), Minhui Zhu, Ping Li, School of Science, Xi'an Polytechnic University, No. 19 Jinhua South Road, Xi'an 710048, Shaanxi, P. R. China, e-mail: `muquanwu@163.com`, `xiao-yu123@126.com`, `liping7487@163.com`.