

LIBERA AND HILBERT MATRIX OPERATOR
ON LOGARITHMICALLY WEIGHTED BERGMAN,
BLOCH AND HARDY-BLOCH SPACES

BOBAN KARAPETROVIĆ, Belgrade

Received October 23, 2016. First published March 6, 2018.

Abstract. We show that if $\alpha > 1$, then the logarithmically weighted Bergman space $A_{\log^\alpha}^2$ is mapped by the Libera operator \mathcal{L} into the space $A_{\log^{\alpha-1}}^2$, while if $\alpha > 2$ and $0 < \varepsilon \leq \alpha - 2$, then the Hilbert matrix operator H maps $A_{\log^\alpha}^2$ into $A_{\log^{\alpha-2-\varepsilon}}^2$.

We show that the Libera operator \mathcal{L} maps the logarithmically weighted Bloch space $\mathcal{B}_{\log^\alpha}$, $\alpha \in \mathbb{R}$, into itself, while H maps $\mathcal{B}_{\log^\alpha}$ into $\mathcal{B}_{\log^{\alpha+1}}$.

In Pavlović's paper (2016) it is shown that \mathcal{L} maps the logarithmically weighted Hardy-Bloch space $\mathcal{B}_{\log^\alpha}^1$, $\alpha > 0$, into $\mathcal{B}_{\log^{\alpha-1}}^1$. We show that this result is sharp. We also show that H maps $\mathcal{B}_{\log^\alpha}^1$, $\alpha \geq 0$, into $\mathcal{B}_{\log^{\alpha-1}}^1$ and that this result is sharp also.

Keywords: Libera operator; Hilbert matrix operator; Hardy space; Bergman space; Bloch space; Hardy-Bloch space

MSC 2010: 47B38, 47G10, 30H25

1. INTRODUCTION

We consider the action of the Libera and Hilbert matrix operators on logarithmically weighted Bergman, Bloch and Hardy-Bloch spaces.

We show that if $\alpha > 1$, then the logarithmically weighted Bergman space $A_{\log^\alpha}^2$ is mapped by the Libera operator \mathcal{L} into the space $A_{\log^{\alpha-1}}^2$. In [4] it is shown that if $f \in A_{\log^\alpha}^2$, where $\alpha > 3$, then $Hf \in A^2$. Here H is the Hilbert matrix operator. Also, in [1] it is shown that $H: A_{\log^\alpha}^2 \rightarrow A^2$ for $\alpha > 2$. We improve this result by showing that if $\alpha > 2$ and $0 < \varepsilon \leq \alpha - 2$, then H is well defined on $A_{\log^\alpha}^2$ and maps this space into $A_{\log^{\alpha-2-\varepsilon}}^2$.

The research has been supported by NTR Serbia, Project ON174032.

We show that the Libera operator \mathcal{L} maps the logarithmically weighted Bloch space $\mathcal{B}_{\log^\alpha}$, $\alpha \in \mathbb{R}$, into itself, while H maps $\mathcal{B}_{\log^\alpha}$ into $\mathcal{B}_{\log^{\alpha+1}}$.

In [8] it is shown that \mathcal{L} maps the logarithmically weighted Hardy-Bloch space $\mathcal{B}_{\log^\alpha}^1$, $\alpha > 0$, into $\mathcal{B}_{\log^{\alpha-1}}^1$. We note that this result is sharp. Our main results are given in Theorem 5.3. Among other things, we show that H maps $\mathcal{B}_{\log^\alpha}^1$, $\alpha \geq 0$, into $\mathcal{B}_{\log^{\alpha-1}}^1$ and that this result is sharp.

The definitions of logarithmically weighted Bergman, Bloch and Hardy-Bloch spaces will be given in Sections 3, 4 and 5, respectively.

For $0 < p \leq \infty$, Hardy space H^p is the space of all functions f holomorphic in the unit disk $\mathbb{D} = \{z \in \mathbb{C}: |z| < 1\}$ for which

$$\|f\|_{H^p} = \|f\|_p = \sup_{0 \leq r < 1} M_p(r, f) < \infty,$$

where

$$M_p(r, f) = \left(\frac{1}{2\pi} \int_0^{2\pi} |f(re^{it})|^p dt \right)^{1/p} \quad \text{if } 0 < p < \infty,$$

and

$$M_\infty(r, f) = \sup_{0 \leq t < 2\pi} |f(re^{it})|.$$

The Lebesgue measure on \mathbb{D} will be denoted by A and will be normalized so as to have $A(\mathbb{D}) = 1$. That is,

$$dA(z) = \frac{1}{\pi} dx dy = \frac{1}{\pi} r dr dt, \quad \text{where } z = x + iy = re^{it}.$$

The Bergman space A^p , $0 < p < \infty$, is the space of holomorphic functions in $L^p(\mathbb{D}, dA)$, that is,

$$A^p = \left\{ f \in \mathcal{H}(\mathbb{D}) : \|f\|_{A^p}^p = \int_{\mathbb{D}} |f(z)|^p dA(z) < \infty \right\}.$$

A function f holomorphic in the unit disk \mathbb{D} belongs to the Hardy-Bloch space $\mathcal{B}_0^{p,q}$, $0 < p \leq \infty$, $0 < q \leq \infty$ (notation from [7]) if

$$\int_0^1 M_p^q(r, f')(1-r)^{q-1} dr < \infty \quad \text{for } 0 < q < \infty,$$

and

$$\sup_{0 \leq r < 1} (1-r)M_p(r, f') < \infty \quad \text{for } q = \infty.$$

Let $\mathcal{H}(\mathbb{D})$ denote the space of all functions holomorphic in the unit disk \mathbb{D} of the complex plane endowed with the topology of uniform convergence on compact

subsets of \mathbb{D} . The dual of $\mathcal{H}(\mathbb{D})$ is equal to $\mathcal{H}(\overline{\mathbb{D}})$, where $g \in \mathcal{H}(\overline{\mathbb{D}})$ means that g is holomorphic in a neighborhood of $\overline{\mathbb{D}}$ (depending on g). The duality pairing is given by

$$\langle f, g \rangle = \sum_{n=0}^{\infty} \widehat{f}(n) \overline{\widehat{g}(n)},$$

where $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n) z^n \in \mathcal{H}(\mathbb{D})$ and $g(z) = \sum_{n=0}^{\infty} \widehat{g}(n) z^n \in \mathcal{H}(\overline{\mathbb{D}})$.

It is easy to see that the Libera operator defined by

$$\overline{\mathcal{L}}g(z) = \sum_{n=0}^{\infty} \left(\sum_{k=n}^{\infty} \frac{\widehat{g}(k)}{k+1} \right) z^n = \int_0^1 g(t + (1-t)z) dt,$$

$$g(z) = \sum_{n=0}^{\infty} \widehat{g}(n) z^n \in \mathcal{H}(\overline{\mathbb{D}})$$

maps $\mathcal{H}(\overline{\mathbb{D}})$ into $\mathcal{H}(\overline{\mathbb{D}})$.

We denote by \mathcal{L} the operator

$$\begin{aligned} \mathcal{L}g(z) &= \int_0^1 g(t + (1-t)z) dt, \\ g(z) &= \sum_{n=0}^{\infty} \widehat{g}(n) z^n \in \mathcal{H}(\mathbb{D}), \end{aligned}$$

whenever the integral converges uniformly on compact subsets of \mathbb{D} . Uniform convergence means that the limit

$$\lim_{r \rightarrow 1^-} \int_0^r g(t + (1-t)z) dt$$

is uniform with respect to z in any compact subset of \mathbb{D} . This hypothesis guarantees that $\mathcal{L}g$ is a holomorphic function in \mathbb{D} . We call \mathcal{L} also the Libera operator since $\mathcal{L} = \overline{\mathcal{L}}$ on $\mathcal{H}(\overline{\mathbb{D}})$.

The Hilbert matrix is an infinite matrix $H = [h_{n,k}]_{n,k=0}^{\infty}$ whose entries are $h_{n,k} = 1/(n+k+1)$ for all nonnegative integers n and k . It can be viewed as an operator on spaces of holomorphic functions by its action on their Taylor coefficients. If

$$f(z) = \sum_{n=0}^{\infty} \widehat{f}(n) z^n$$

is a holomorphic function in the unit disk \mathbb{D} , then we define the transformation H by

$$Hf(z) = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} h_{n,k} \widehat{f}(k) z^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\widehat{f}(k)}{n+k+1} z^n.$$

It is possible to write Hf , $f \in H^p$, $1 \leq p \leq \infty$, in an integral form, which is quite convenient for analyzing the operator. More specifically, by looking at the Taylor series expansion of the function f , we have the following integral representation:

$$Hf(z) = \int_0^1 \frac{f(t)}{1-tz} dt.$$

It is well known that the Libera operator \mathcal{L} acts as a bounded operator from H^p into H^p if and only if $1 < p \leq \infty$ and that \mathcal{L} acts as a bounded operator from A^p into A^p if and only if $2 < p < \infty$ (see [2], [6]). On the other hand, it is well known that the Hilbert matrix operator H acts as a bounded operator from H^p into H^p if and only if $1 < p < \infty$ and that H acts as a bounded operator from A^p into A^p if and only if $2 < p < \infty$ (see [3]).

2. SOME PRELIMINARY RESULTS

In this section we shall collect some results which will be needed in our work. We start with one useful result.

Sublemma 2.1. *Let $\alpha \in \mathbb{R}$ and $a \geq 2$. Then*

$$\int_{\log a}^{\infty} t^{\alpha} e^{-t} dt \leq C_{\alpha} \frac{\log^{\alpha} a}{a},$$

where C_{α} is a constant independent of a .

Proof. (1) Case $\alpha \leq 0$.

$$\int_{\log a}^{\infty} t^{\alpha} e^{-t} dt \leq \log^{\alpha} a \int_{\log a}^{\infty} e^{-t} dt = \frac{\log^{\alpha} a}{a}.$$

(2) Case $\alpha > 0$. In this case, partial integration gives

$$\begin{aligned} \int_{\log a}^{\infty} t^{\alpha} e^{-t} dt &= \frac{\log^{\alpha} a}{a} + \alpha \int_{\log a}^{\infty} t^{\alpha-1} e^{-t} dt \\ &= \frac{\log^{\alpha} a}{a} + \alpha \frac{\log^{\alpha-1} a}{a} + \alpha(\alpha-1) \int_{\log a}^{\infty} t^{\alpha-2} e^{-t} dt \\ &\leq C_{\alpha} \frac{\log^{\alpha} a}{a} + \alpha(\alpha-1) \int_{\log a}^{\infty} t^{\alpha-2} e^{-t} dt. \end{aligned}$$

Continuing on this way, we find that

$$\begin{aligned}
\int_{\log a}^{\infty} t^{\alpha} e^{-t} dt &\leq C_{\alpha} \frac{\log^{\alpha} a}{a} + \alpha(\alpha-1) \dots (\alpha - [\alpha]) \int_{\log a}^{\infty} t^{\alpha - [\alpha] - 1} e^{-t} dt \\
&\leq C_{\alpha} \frac{\log^{\alpha} a}{a} + \alpha(\alpha-1) \dots (\alpha - [\alpha]) \frac{\log^{\alpha - [\alpha] - 1} a}{a} \\
&\leq C_{\alpha} \frac{\log^{\alpha} a}{a},
\end{aligned}$$

where $[\alpha]$ is the largest integer less than or equal to α . □

Consequently, we get the following result.

Lemma 2.2. *Let $\alpha \in \mathbb{R}$ and let n be a nonnegative integer. Then*

$$\int_0^1 r^n \log^{\alpha} \frac{2}{1-r} dr \asymp \frac{\log^{\alpha}(n+2)}{n+1},$$

where the corresponding constant is independent of n , i.e., there is a constant C_{α} independent of n such that

$$\frac{1}{C_{\alpha}} \frac{\log^{\alpha}(n+2)}{n+1} \leq \int_0^1 r^n \log^{\alpha} \frac{2}{1-r} dr \leq C_{\alpha} \frac{\log^{\alpha}(n+2)}{n+1}.$$

Proof. (1) Case $\alpha \geq 0$. First, we find that

$$\begin{aligned}
\int_0^1 r^n \log^{\alpha} \frac{2}{1-r} dr &\geq \int_{1-1/(n+1)}^1 r^n \log^{\alpha} \frac{2}{1-r} dr \\
&\geq \log^{\alpha}(n+2) \int_{1-1/(n+1)}^1 r^n dr \\
&= \frac{\log^{\alpha}(n+2)}{n+1} \left(1 - \left(1 - \frac{1}{n+1}\right)^{n+1}\right) \\
&\geq C \frac{\log^{\alpha}(n+2)}{n+1}.
\end{aligned}$$

On the other hand, by using Sublemma 2.1, we have that

$$\begin{aligned}
\int_{1-1/(n+1)}^1 r^n \log^{\alpha} \frac{2}{1-r} dr &\leq \int_{1-1/(n+1)}^1 \log^{\alpha} \frac{2}{1-r} dr \\
&= 2 \int_{\log(2n+2)}^{\infty} t^{\alpha} e^{-t} dt \\
&\leq 2 \int_{\log(n+2)}^{\infty} t^{\alpha} e^{-t} dt \\
&\leq C_{\alpha} \frac{\log^{\alpha}(n+2)}{n+1},
\end{aligned}$$

and

$$\begin{aligned} \int_0^{1-1/(n+1)} r^n \log^\alpha \frac{2}{1-r} dr &\leq C_\alpha \log^\alpha(n+2) \int_0^{1-1/(n+1)} r^n dr \\ &= C_\alpha \frac{\log^\alpha(n+2)}{n+1} \left(1 - \frac{1}{n+1}\right)^{n+1} \\ &\leq C_\alpha \frac{\log^\alpha(n+2)}{n+1}. \end{aligned}$$

Therefore,

$$\int_0^1 r^n \log^\alpha \frac{2}{1-r} dr \leq C_\alpha \frac{\log^\alpha(n+2)}{n+1}.$$

(2) Case $\alpha < 0$. Let $\varphi(r) = r \log^\alpha(2/r)$, $0 < r \leq 1$. Then φ is a nonnegative, increasing function on the interval $(0, 1]$ and

$$t^{1-2\alpha} \varphi(r) \leq \varphi(tr) \leq t \varphi(r)$$

for all $0 < t < 1$. By using Lemma 4.1 in [5], we find that

$$\int_0^1 r^n \frac{\varphi(1-r)}{1-r} dr \asymp \varphi\left(\frac{1}{n+1}\right).$$

Hence,

$$\int_0^1 r^n \log^\alpha \frac{2}{1-r} dr \asymp \frac{\log^\alpha(n+2)}{n+1}.$$

□

The following auxiliary result will be useful.

Theorem 2.3. (a) For every real α , the Taylor coefficients $\widehat{F}(n)$ of the function

$$F(z) = \frac{1}{1-z} \log^\alpha \frac{2}{1-z}$$

have the property

$$\widehat{F}(n) \asymp \log^\alpha(n+2),$$

where the corresponding constant is independent of n .

(b) For every real α , the Taylor coefficients $\widehat{G}(n)$ of the function

$$G(z) = \log^\alpha \frac{2}{1-z}$$

have the property

$$\widehat{G}(n) \asymp \frac{\log^{\alpha-1}(n+2)}{n+1},$$

where the corresponding constant is independent of n .

This theorem is a consequence of Theorem 2.31 on page 192 of the classical monograph [10], hence we omit its proof. Now, we are ready to prove our next result.

Lemma 2.4. *Let $\alpha \in \mathbb{R}$ and let k be a nonnegative integer. Then*

$$\sum_{n=0}^{\infty} \frac{\log^{\alpha}(n+2)}{(n+1)(n+k+1)} \asymp \frac{\log^{\alpha+1}(k+2)}{k+1},$$

where the corresponding constant is independent of k , i.e., there is a constant C_{α} independent of k such that

$$\frac{1}{C_{\alpha}} \frac{\log^{\alpha+1}(k+2)}{k+1} \leq \sum_{n=0}^{\infty} \frac{\log^{\alpha}(n+2)}{(n+1)(n+k+1)} \leq C_{\alpha} \frac{\log^{\alpha+1}(k+2)}{k+1}.$$

Proof. By using Lemma 2.2 and Theorem 2.3 (b), we find that

$$\begin{aligned} \frac{\log^{\alpha+1}(k+2)}{k+1} &\asymp \int_0^1 r^k \log^{\alpha+1} \frac{2}{1-r} dr \\ &\asymp \int_0^1 r^k \sum_{n=0}^{\infty} \frac{\log^{\alpha}(n+2)}{n+1} r^n dr \\ &= \sum_{n=0}^{\infty} \frac{\log^{\alpha}(n+2)}{n+1} \int_0^1 r^{n+k} dr \\ &= \sum_{n=0}^{\infty} \frac{\log^{\alpha}(n+2)}{(n+1)(n+k+1)}, \end{aligned}$$

where the corresponding constant is independent of k . □

3. LIBERA AND HILBERT MATRIX OPERATOR ON LOGARITHMICALLY WEIGHTED BERGMAN SPACES

For $\alpha \in \mathbb{R}$ we define the logarithmically weighted Bergman spaces $A_{\log^{\alpha}}^2$ as follows:

$$A_{\log^{\alpha}}^2 = \left\{ f \in \mathcal{H}(\mathbb{D}) : \|f\|_{A_{\log^{\alpha}}^2}^2 = \int_{\mathbb{D}} |f(z)|^2 \log^{\alpha} \frac{2}{1-|z|^2} dA(z) < \infty \right\}.$$

Note that $A_{\log^{\alpha}}^2 \subset A^2$ for $\alpha > 0$ and $A_{\log^0}^2 = A^2$.

Let $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n) z^n \in A_{\log^{\alpha}}^2$. By using Parseval's formula and Lemma 2.2, we find that

$$\|f\|_{A_{\log^{\alpha}}^2}^2 = \sum_{n=0}^{\infty} |\widehat{f}(n)|^2 \int_0^1 r^n \log^{\alpha} \frac{2}{1-r} dr \asymp \sum_{n=0}^{\infty} |\widehat{f}(n)|^2 \frac{\log^{\alpha}(n+2)}{n+1},$$

where the corresponding constant is independent of function f , i.e., there is a constant C independent of function f such that

$$\frac{1}{C} \|f\|_{A_{\log^\alpha}^2}^2 \leq \sum_{n=0}^{\infty} |\widehat{f}(n)|^2 \frac{\log^\alpha(n+2)}{n+1} \leq C \|f\|_{A_{\log^\alpha}^2}^2.$$

3.1. Libera operator on logarithmically weighted Bergman spaces. Our next result describes the action of the Libera operator \mathcal{L} on the logarithmically weighted Bergman space $A_{\log^\alpha}^2$ for $\alpha > 1$.

Theorem 3.1. *If $\alpha > 1$, then the operator \mathcal{L} is well defined on $A_{\log^\alpha}^2$ and maps this space into $A_{\log^{\alpha-1}}^2$.*

Proof. Let $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n) z^n \in A_{\log^\alpha}^2$. Then, by using the Cauchy-Schwarz inequality, we find that

$$\sum_{n=0}^{\infty} \frac{|\widehat{f}(n)|}{n+1} \leq \left(\sum_{n=0}^{\infty} |\widehat{f}(n)|^2 \frac{\log^\alpha(n+2)}{n+1} \right)^{1/2} \left(\sum_{n=0}^{\infty} \frac{1}{(n+1) \log^\alpha(n+2)} \right)^{1/2} < \infty,$$

because $\alpha > 1$. From this, we get that if $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n) z^n \in A_{\log^\alpha}^2$, then $\sum_{n=0}^{\infty} |\widehat{f}(n)| \times (n+1)^{-1} < \infty$. Hence, the operator \mathcal{L} is well defined on $A_{\log^\alpha}^2$. Using inequality (59) from [6], we find that

$$r M_2^2(r, \mathcal{L}f) \leq C(1-r)^{-1} \int_r^1 M_2^2(s, f) ds$$

for all $0 \leq r < 1$. Therefore,

$$\begin{aligned} \int_{\mathbb{D}} |\mathcal{L}f(z)|^2 \log^{\alpha-1} \frac{2}{1-|z|^2} dA(z) &= 2 \int_0^1 r M_2^2(r, \mathcal{L}f) \log^{\alpha-1} \frac{2}{1-r^2} dr \\ &\leq C \int_0^1 \frac{1}{1-r} \int_r^1 M_2^2(s, f) ds \log^{\alpha-1} \frac{2}{1-r^2} dr \\ &= C \int_0^1 M_2^2(s, f) \int_0^s \frac{1}{1-r} \log^{\alpha-1} \frac{2}{1-r^2} dr ds \\ &\leq C \int_0^1 M_2^2(s, f) \int_0^s \frac{1}{1-r} \log^{\alpha-1} \frac{2}{1-r} dr ds \\ &= C \int_0^1 M_2^2(s, f) \left(\log^\alpha \frac{2}{1-s} - \log^\alpha 2 \right) ds \\ &\leq C \int_0^1 M_2^2(s, f) \log^\alpha \frac{2}{1-s} ds \end{aligned}$$

$$\begin{aligned}
&= C \int_0^1 u M_2^2(u^2, f) \log^\alpha \frac{2}{1-u^2} du \\
&\leq C \int_0^1 u M_2^2(u, f) \log^\alpha \frac{2}{1-u^2} du \\
&= C \int_{\mathbb{D}} |f(z)|^2 \log^\alpha \frac{2}{1-|z|^2} dA(z) < \infty.
\end{aligned}$$

Hence, $\mathcal{L}f \in A_{\log^{\alpha-1}}^2$. □

3.2. Hilbert matrix operator on logarithmically weighted Bergman spaces. In [4] it is shown that if $f \in A_{\log^\alpha}^2$, where $\alpha > 3$, then $Hf \in A^2$. Also, in [1] it is shown that $H: A_{\log^\alpha}^2 \rightarrow A^2$ for $\alpha > 2$. Our next theorem improves this result.

Theorem 3.2. *If $\alpha > 2$ and $0 < \varepsilon \leq \alpha - 2$, then H is well defined on $A_{\log^\alpha}^2$ and maps this space into $A_{\log^{\alpha-2-\varepsilon}}^2$.*

Proof. For $\alpha > 2$, we have that if $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n)z^n \in A_{\log^\alpha}^2$, then we get $\sum_{n=0}^{\infty} |\widehat{f}(n)|/(n+1) < \infty$. Therefore, the operator H is well defined on $A_{\log^\alpha}^2$. On the other hand, if $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n)z^n \in A_{\log^\alpha}^2$, then by using the Cauchy-Schwarz inequality and Lemma 2.4, we find that

$$\begin{aligned}
\|Hf\|_{A_{\log^{\alpha-2-\varepsilon}}^2}^2 &\asymp \sum_{n=0}^{\infty} |\widehat{Hf}(n)|^2 \frac{\log^{\alpha-2-\varepsilon}(n+2)}{n+1} \\
&= \sum_{n=0}^{\infty} \left| \sum_{k=0}^{\infty} \frac{\widehat{f}(k)}{n+k+1} \right|^2 \frac{\log^{\alpha-2-\varepsilon}(n+2)}{n+1} \\
&\leq \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{|\widehat{f}(k)|^2 \log^\alpha(k+2)}{n+k+1} \\
&\quad \times \sum_{k=0}^{\infty} \frac{1}{(n+k+1) \log^\alpha(k+2)} \frac{\log^{\alpha-2-\varepsilon}(n+2)}{n+1} \\
&\leq \sum_{k=0}^{\infty} |\widehat{f}(k)|^2 \frac{\log^\alpha(k+2)}{k+1} \sum_{k=0}^{\infty} \frac{1}{\log^\alpha(k+2)} \sum_{n=0}^{\infty} \frac{\log^{\alpha-2-\varepsilon}(n+2)}{(n+1)(n+k+1)} \\
&\leq C \|f\|_{A_{\log^\alpha}^2}^2 \sum_{k=0}^{\infty} \frac{1}{\log^\alpha(k+2)} \frac{\log^{\alpha-1-\varepsilon}(k+2)}{k+1} \\
&= C \|f\|_{A_{\log^\alpha}^2}^2 \sum_{k=0}^{\infty} \frac{1}{(k+1) \log^{1+\varepsilon}(k+2)} < \infty.
\end{aligned}$$

Therefore, $Hf \in A_{\log^{\alpha-2-\varepsilon}}^2$. □

We note that for $\alpha \in (1, 2]$ the operator H is well defined on $A_{\log^\alpha}^2$. We do not know whether Theorem 3.2 holds in this case. A natural question is: Does Theorem 3.2 hold for $\varepsilon = 0$?

4. LIBERA AND HILBERT MATRIX OPERATOR ON LOGARITHMICALLY WEIGHTED BLOCH SPACES

For $\alpha \in \mathbb{R}$ we define the logarithmically weighted Bloch spaces $\mathcal{B}_{\log^\alpha}$ as follows,

$$\mathcal{B}_{\log^\alpha} = \left\{ f \in \mathcal{H}(\mathbb{D}) : |f'(z)|(1 - |z|) = \mathcal{O}\left(\log^\alpha \frac{2}{1 - |z|}\right) \right\}.$$

The norm in the space $\mathcal{B}_{\log^\alpha}$ is defined by

$$\|f\|_{\mathcal{B}_{\log^\alpha}} = |f(0)| + \sup_{z \in \mathbb{D}} |f'(z)|(1 - |z|) \log^{-\alpha} \frac{2}{1 - |z|}.$$

Note that \mathcal{B}_{\log^0} is the Bloch space $\mathcal{B}_0^{\infty, \infty} = \mathcal{B}$.

4.1. Libera operator on logarithmically weighted Bloch spaces. Now, we have the following theorem.

Theorem 4.1. *Let $\alpha \in \mathbb{R}$. Then \mathcal{L} is well defined on $\mathcal{B}_{\log^\alpha}$ and maps this space into $\mathcal{B}_{\log^\alpha}$.*

Proof. By Theorem 2.1 (a) in [8], we have that if $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n)z^n \in \mathcal{B}_{\log^\alpha}$, then $\sum_{n=0}^{\infty} |\widehat{f}(n)|/(n+1) < \infty$. Therefore, the Libera operator \mathcal{L} is well defined on $\mathcal{B}_{\log^\alpha}$.

By using Lemma 22 from [6], for $\nu = 1$ and $p = \infty$, we find that

$$M_\infty(r, (\mathcal{L}f)') \leq (1 - r)^{-2} \int_r^1 (1 - s) M_\infty(s, f') ds$$

for all $0 \leq r < 1$. Then, by using Sublemma 2.1, we have that

$$\begin{aligned} (1 - r) M_\infty(r, (\mathcal{L}f)') &\leq \frac{1}{1 - r} \int_r^1 (1 - s) M_\infty(s, f') ds \\ &\leq C \frac{1}{1 - r} \int_r^1 \log^\alpha \frac{2}{1 - s} ds \\ &= C \frac{1}{1 - r} \int_{\log(2/(1-r))}^{\infty} t^\alpha e^{-t} dt \\ &\leq C \log^\alpha \frac{2}{1 - r}. \end{aligned}$$

Hence, we obtain $\mathcal{L}f \in \mathcal{B}_{\log^\alpha}$. □

4.2. Hilbert matrix operator on logarithmically weighted Bloch spaces.

Our next result describes the action of the Hilbert matrix operator H on the logarithmically weighted Bloch space $\mathcal{B}_{\log^\alpha}$ for $\alpha \in \mathbb{R}$. We improve results given in Proposition 5.1 and Proposition 5.2 in [4].

Theorem 4.2. *Let $\alpha \in \mathbb{R}$. Then H is well defined on $\mathcal{B}_{\log^\alpha}$ and maps this space into $\mathcal{B}_{\log^{\alpha+1}}$.*

Proof. By Theorem 2.1 (a) in [8], we have that if $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n)z^n \in \mathcal{B}_{\log^\alpha}$, then $\sum_{n=0}^{\infty} |\widehat{f}(n)|/(n+1) < \infty$. Hence, the Hilbert matrix operator H is well defined on $\mathcal{B}_{\log^\alpha}$.

Now, let $f \in \mathcal{B}_{\log^\alpha}$, where without loss of generality, we can additionally assume that $f(0) = 0$. Then, by Lemma 4.2.8 in [9], we can write

$$f(z) = \int_{\mathbb{D}} \frac{f'(w)(1-|w|^2)}{\overline{w}(1-\overline{w}z)^2} dA(w),$$

for all $z \in \mathbb{D}$. Also, we have that

$$\int_{\mathbb{D}} \frac{z^k |z|^{2n}}{1-\overline{z}} dA(z) = \frac{1}{n+k+1},$$

for all nonnegative integers n and k . Therefore,

$$\int_{\mathbb{D}} f(z) \frac{|z|^{2n}}{1-\overline{z}} dA(z) = \sum_{k=0}^{\infty} \frac{\widehat{f}(k)}{n+k+1} = \widehat{Hf}(n).$$

Consequently,

$$\begin{aligned} |\widehat{Hf}(n)| &= \left| \int_{\mathbb{D}} f(z) \frac{|z|^{2n}}{1-\overline{z}} dA(z) \right| \\ &= \left| \int_{\mathbb{D}} \frac{f'(w)(1-|w|^2)}{\overline{w}} \int_{\mathbb{D}} \frac{|z|^{2n}}{(1-\overline{w}z)^2(1-\overline{z})} dA(z) dA(w) \right| \\ &= \frac{1}{\pi} \left| \int_{\mathbb{D}} \frac{f'(w)(1-|w|^2)}{\overline{w}} \int_0^1 r^{2n+1} \int_0^{2\pi} \frac{1}{(1-\overline{w}re^{i\theta})^2(1-re^{-i\theta})} d\theta dr dA(w) \right| \\ &= \frac{1}{\pi} \left| \int_{\mathbb{D}} \frac{f'(w)(1-|w|^2)}{\overline{w}} \int_0^1 r^{2n+1} \frac{2\pi}{(1-r^2\overline{w})^2} dr dA(w) \right| \\ &\leq C \int_{\mathbb{D}} \frac{|f'(w)|(1-|w|)}{|w|} \int_0^1 \frac{r^{2n+1}}{|1-r^2\overline{w}|^2} dr dA(w) \\ &\leq C \int_{\mathbb{D}} \frac{\log^\alpha \frac{2}{1-|w|}}{|w|} \int_0^1 \frac{r^{2n+1}}{|1-r^2\overline{w}|^2} dr dA(w) \end{aligned}$$

$$\begin{aligned}
&= C \int_0^1 r^{2n+1} \int_{\mathbb{D}} \frac{\log^\alpha \frac{2}{1-|w|}}{|w||1-r^2w|^2} dA(w) dr \\
&= C \int_0^1 r^{2n+1} \int_0^1 \log^\alpha \frac{2}{1-\varrho} \int_0^{2\pi} \frac{1}{|1-r^2\varrho e^{i\theta}|^2} d\theta d\varrho dr \\
&\leq C \int_0^1 r^{2n+1} \int_0^1 \frac{1}{1-r^2\varrho} \log^\alpha \frac{2}{1-\varrho} d\varrho dr \\
&= C \int_0^1 \log^\alpha \frac{2}{1-\varrho} \int_0^1 \frac{r^n}{1-r\varrho} dr d\varrho.
\end{aligned}$$

On the other hand, we find that

$$\int_0^1 \frac{r^n}{1-r\varrho} dr = \sum_{k=0}^{\infty} \frac{\varrho^k}{n+k+1}.$$

Hence, by using Lemma 2.2 and Lemma 2.4, we obtain

$$\begin{aligned}
|\widehat{Hf}(n)| &\leq C \sum_{k=0}^{\infty} \frac{1}{n+k+1} \int_0^1 \varrho^k \log^\alpha \frac{2}{1-\varrho} d\varrho \\
&\leq C \sum_{k=0}^{\infty} \frac{\log^\alpha(k+2)}{(k+1)(n+k+1)} \\
&\leq C \frac{\log^{\alpha+1}(n+2)}{n+1}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
|(Hf)'(z)| &= \left| \sum_{n=1}^{\infty} n \widehat{Hf}(n) z^{n-1} \right| \\
&\leq \sum_{n=1}^{\infty} n |\widehat{Hf}(n)| |z|^{n-1} \\
&\leq C \sum_{n=1}^{\infty} \frac{n}{n+1} \log^{\alpha+1}(n+2) |z|^{n-1} \\
&\leq C \sum_{n=0}^{\infty} \log^{\alpha+1}(n+3) |z|^n \\
&\leq C \sum_{n=0}^{\infty} \log^{\alpha+1}(n+2) |z|^n.
\end{aligned}$$

By using Theorem 2.3 (a), we find that

$$\frac{1}{1-|z|} \log^{\alpha+1} \frac{2}{1-|z|} \asymp \sum_{n=0}^{\infty} \log^{\alpha+1}(n+2) |z|^n.$$

Finally,

$$\begin{aligned} |(Hf)'(z)| &\leq C \sum_{n=0}^{\infty} \log^{\alpha+1}(n+2) |z|^n \\ &\leq C \frac{1}{1-|z|} \log^{\alpha+1} \frac{2}{1-|z|}. \end{aligned}$$

Hence, $Hf \in \mathcal{B}_{\log^{\alpha+1}}$. □

5. LIBERA AND HILBERT MATRIX OPERATOR ON LOGARITHMICALLY WEIGHTED HARDY-BLOCH SPACES

For $\alpha \in \mathbb{R}$ we define the logarithmically weighted Hardy-Bloch spaces $\mathcal{B}_{\log^{\alpha}}^1$ in the following way:

$$\mathcal{B}_{\log^{\alpha}}^1 = \left\{ f \in \mathcal{H}(\mathbb{D}) : \|f\|_{\mathcal{B}_{\log^{\alpha}}^1} = |f(0)| + \int_{\mathbb{D}} |f'(z)| \log^{\alpha} \frac{2}{1-|z|} dA(z) < \infty \right\}.$$

For $\alpha = 0$, $\mathcal{B}_{\log^0}^1$ is the Hardy-Bloch space $\mathcal{B}_0^{1,1}$ (notation from [7]). We note that if $\alpha \geq 0$, then $\mathcal{B}_{\log^{\alpha}}^1 \subseteq \mathcal{B}_0^{1,1} \subseteq H^1$ and if $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n) z^n \in H^1$, then $\sum_{n=0}^{\infty} |\widehat{f}(n)|/(n+1) < \infty$.

5.1. Libera operator on logarithmically weighted Hardy-Bloch spaces.

Action of the Libera operator on the logarithmically weighted Hardy-Bloch spaces has been considered in [8] and the following two theorems are proved.

Theorem 5.1 ([8]). *Let $\alpha \geq -1$ and let $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n) z^n \in \mathcal{H}(\mathbb{D})$, where $\widehat{f}(n) \downarrow 0$ as $n \rightarrow \infty$. Then*

$$f \in \mathcal{B}_{\log^{\alpha}}^1 \quad \text{if and only if} \quad \sum_{n=0}^{\infty} \widehat{f}(n) \frac{\log^{\alpha}(n+2)}{n+1} < \infty.$$

Moreover, there is a constant C independent of the function f , such that

$$\frac{1}{C} \|f\|_{\mathcal{B}_{\log^{\alpha}}^1} \leq \sum_{n=0}^{\infty} \widehat{f}(n) \frac{\log^{\alpha}(n+2)}{n+1} \leq C \|f\|_{\mathcal{B}_{\log^{\alpha}}^1}.$$

Theorem 5.2 ([8]). *Let $\alpha > 0$.*

- (a) *Then \mathcal{L} is well defined on $\mathcal{B}_{\log^\alpha}^1$ and maps this space into $\mathcal{B}_{\log^{\alpha-1}}^1$.*
- (b) *If $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n)z^n$, where $\widehat{f}(n) \downarrow 0$ as $n \rightarrow \infty$ and $\sum_{n=0}^{\infty} \widehat{f}(n)/(n+1) < \infty$, then $\mathcal{L}f \in \mathcal{B}_{\log^{\alpha-1}}^1$ implies $f \in \mathcal{B}_{\log^\alpha}^1$.*
- (c) *If $\alpha < 0$, then $\overline{\mathcal{L}}$ cannot be extended to a continuous operator from $\mathcal{B}_{\log^\alpha}^1$ to $\mathcal{H}(\mathbb{D})$.*

For (a) see Theorem 2.3 in [8]. Item (b) follows from Theorem 1.1 and Theorem 1.2 in [8]. For (c) see Theorem 2.1 (c) in [8].

5.2. Hilbert matrix operator on logarithmically weighted Hardy-Bloch spaces. Now we are ready to state the main theorem of this section.

Theorem 5.3. *Let $\alpha \geq 0$.*

- (a) *Then H is well defined on $\mathcal{B}_{\log^\alpha}^1$ and maps this space into $\mathcal{B}_{\log^{\alpha-1}}^1$.*
- (b) *If $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n)z^n$, where $\widehat{f}(n) \downarrow 0$ as $n \rightarrow \infty$ and $\sum_{n=0}^{\infty} \widehat{f}(n)/(n+1) < \infty$, then $Hf \in \mathcal{B}_{\log^{\alpha-1}}^1$ implies $f \in \mathcal{B}_{\log^\alpha}^1$.*
- (c) *The result in (a) is sharp in the sense that for any $\varepsilon > 0$ there exists $f \in \mathcal{B}_{\log^\alpha}^1$ such that $Hf \notin \mathcal{B}_{\log^{\alpha-1+\varepsilon}}^1$.*
- (d) *If $\alpha < 0$, then H cannot be extended to a continuous operator from $\mathcal{B}_{\log^\alpha}^1$ to $\mathcal{H}(\mathbb{D})$.*

Proof. (a) By Theorem 2.1 (b) in [8], we have that if $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n)z^n \in \mathcal{B}_{\log^\alpha}^1$, then $\sum_{n=0}^{\infty} |\widehat{f}(n)|/(n+1) < \infty$. Therefore, the operator H is well defined on $\mathcal{B}_{\log^\alpha}^1$.

Now, let $f \in \mathcal{B}_{\log^\alpha}^1$, where without loss of generality, we can additionally assume that $f(0) = 0$. Then, by Lemma 4.2.8 in [9], we have

$$f(z) = \int_{\mathbb{D}} \frac{f'(w)(1 - |w|^2)}{\overline{w}(1 - \overline{w}z)^2} dA(w),$$

for all $z \in \mathbb{D}$. Let $S = \int_{\mathbb{D}} |(Hf)'(z)| \log^{\alpha-1}(2/(1 - |z|)) dA(z)$. Then, by using the integral representation of the Hilbert matrix operator $Hf(z) = \int_0^1 (f(t)/(1 - tz)) dt$,

we find that

$$\begin{aligned}
S &\leq \int_{\mathbb{D}} \int_0^1 \frac{|f(t)|}{|1-tz|^2} \log^{\alpha-1} \frac{2}{1-|z|} dt dA(z) \\
&= \int_{\mathbb{D}} \int_0^1 \frac{1}{|1-tz|^2} \left| \int_{\mathbb{D}} \frac{f'(w)(1-|w|^2)}{\bar{w}(1-\bar{w}t)^2} dA(w) \right| \log^{\alpha-1} \frac{2}{1-|z|} dt dA(z) \\
&\leq C \int_{\mathbb{D}} \frac{|f'(w)|(1-|w|)}{|w|} \int_{\mathbb{D}} \log^{\alpha-1} \frac{2}{1-|z|} \int_0^1 \frac{dt}{|1-tz|^2|1-tw|^2} dA(z) dA(w) \\
&\leq C \int_{\mathbb{D}} \frac{|f'(w)|(1-|w|)}{|w|} \int_{\mathbb{D}} \frac{1}{|1-zw|^3} \log^{\alpha-1} \frac{2}{1-|z|} dA(z) dA(w) \\
&\leq C \int_{\mathbb{D}} \frac{|f'(w)|}{|w|} \int_{\mathbb{D}} \frac{1}{|1-zw|^2} \log^{\alpha-1} \frac{2}{1-|z|} dA(z) dA(w) \\
&\leq C \int_{\mathbb{D}} \frac{|f'(w)|}{|w|} \int_0^1 \int_0^{2\pi} \frac{d\theta}{|1-re^{i\theta}w|^2} \log^{\alpha-1} \frac{2}{1-r} dr dA(w) \\
&\leq C \int_{\mathbb{D}} \frac{|f'(w)|}{|w|} \int_0^1 \frac{1}{1-r|w|} \log^{\alpha-1} \frac{2}{1-r} dr dA(w).
\end{aligned}$$

On the other hand, by using Lemma 2.2 and Theorem 2.3 (b), we have that

$$\begin{aligned}
\int_0^1 \frac{1}{1-r|w|} \log^{\alpha-1} \frac{2}{1-r} dr &= \sum_{n=0}^{\infty} |w|^n \int_0^1 r^n \log^{\alpha-1} \frac{2}{1-r} dr \\
&\asymp \sum_{n=0}^{\infty} \frac{\log^{\alpha-1}(n+2)}{n+1} |w|^n \\
&\asymp \log^{\alpha} \frac{2}{1-|w|}.
\end{aligned}$$

Consequently,

$$\begin{aligned}
S &\leq C \int_{\mathbb{D}} \frac{|f'(w)|}{|w|} \log^{\alpha} \frac{2}{1-|w|} dA(w) \\
&\leq C \int_{\mathbb{D}} |f'(w)| \log^{\alpha} \frac{2}{1-|w|} dA(w) \\
&< \infty
\end{aligned}$$

and we get that $Hf \in \mathcal{B}_{\log^{\alpha-1}}^1$.

(b) We have $Hf(z) = \sum_{n=0}^{\infty} \widehat{Hf}(n)z^n$, where $\widehat{Hf}(n) = \sum_{k=0}^{\infty} \widehat{f}(k)/(n+k+1) \downarrow 0$ as $n \rightarrow \infty$. Then, by using Theorem 5.1, we find that

$$\|Hf\|_{\mathcal{B}_{\log^{\alpha-1}}^1} \asymp \sum_{n=0}^{\infty} \widehat{Hf}(n) \frac{\log^{\alpha-1}(n+2)}{n+1},$$

where the corresponding constant is independent of f . On the other hand, by using Lemma 2.4, we have that

$$\begin{aligned} \sum_{n=0}^{\infty} \widehat{Hf}(n) \frac{\log^{\alpha-1}(n+2)}{n+1} &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{\widehat{f}(k)}{n+k+1} \frac{\log^{\alpha-1}(n+2)}{n+1} \\ &= \sum_{k=0}^{\infty} \widehat{f}(k) \sum_{n=0}^{\infty} \frac{\log^{\alpha-1}(n+2)}{(n+1)(n+k+1)} \\ &\asymp \sum_{k=0}^{\infty} \widehat{f}(k) \frac{\log^{\alpha}(k+2)}{k+1}. \end{aligned}$$

Therefore,

$$\|Hf\|_{\mathcal{B}_{\log^{\alpha-1}}^1} \asymp \sum_{n=0}^{\infty} \widehat{f}(n) \frac{\log^{\alpha}(n+2)}{n+1},$$

where the corresponding constant is independent of f . Then $\sum_{n=0}^{\infty} \widehat{f}(n) \log^{\alpha}(n+2) \times (n+1)^{-1} < \infty$ and by using Theorem 5.1 we find that $f \in \mathcal{B}_{\log^{\alpha}}^1$.

(c) Let $\varepsilon > 0$ and let $\widehat{f}(n) = (\log^{\alpha+1+\varepsilon/2}(n+2))^{-1}$ for all $n \geq 0$. Then $\sum_{n=0}^{\infty} \widehat{f}(n) \times (n+1)^{-1} < \infty$ and $\widehat{f}(n) \downarrow 0$ as $n \rightarrow \infty$. Also, we find that

$$\sum_{n=0}^{\infty} \widehat{f}(n) \frac{\log^{\alpha}(n+2)}{n+1} < \infty \quad \text{and} \quad \sum_{n=0}^{\infty} \widehat{f}(n) \frac{\log^{\alpha+\varepsilon}(n+2)}{n+1} = \infty.$$

Let $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n) z^n$. Then $f \in \mathcal{B}_{\log^{\alpha}}^1$ by Theorem 5.1 and $Hf \notin \mathcal{B}_{\log^{\alpha-1+\varepsilon}}^1$, because otherwise we would have $\sum_{n=0}^{\infty} \widehat{f}(n) \log^{\alpha+\varepsilon}(n+2)(n+1)^{-1} < \infty$ by part (b) of this theorem. A contradiction.

(d) Since $\mathcal{B}_{\log^{\alpha}}^1 \subset \mathcal{B}_{\log^{\beta}}^1$ for $\beta < \alpha$, we may assume that $-1 < \alpha < 0$. Let

$$f(z) = \sum_{n=0}^{\infty} \frac{z^n}{\log(n+2)}.$$

For every $r \in (0, 1)$ the function $f_r(z) = f(rz)$ belongs to $\mathcal{H}(\overline{\mathbb{D}})$ and by Theorem 5.1, the set $\{f_r : r \in (0, 1)\}$ is bounded in $\mathcal{B}_{\log^{\alpha}}^1$. On the other hand,

$$Hf_r(0) = \sum_{k=0}^{\infty} \frac{r^k}{(k+1) \log(k+2)} \rightarrow \infty \quad \text{as } r \uparrow 1.$$

This contradicts the fact that if a set $X \subset \mathcal{B}_{\log^{\alpha}}^1$ is bounded and H is bounded on $\mathcal{B}_{\log^{\alpha}}^1$, then the set $\{Hf(0) : f \in X\}$ is bounded, because the functional $h \rightarrow h(0)$ is continuous on $\mathcal{H}(\mathbb{D})$. This completes the proof. \square

Remark 5.4. We note that the result stated in Theorem 5.2 (a) is sharp in the sense that for any $\varepsilon > 0$ and $\alpha > 0$ there exists $f \in \mathcal{B}_{\log^\alpha}^1$ such that $\mathcal{L}f \notin \mathcal{B}_{\log^{\alpha-1+\varepsilon}}^1$. As above, we have that

$$f(z) = \sum_{n=0}^{\infty} \frac{z^n}{\log^{\alpha+1+\varepsilon/2}(n+2)} \in \mathcal{B}_{\log^\alpha}^1,$$

while $\mathcal{L}f \notin \mathcal{B}_{\log^{\alpha-1+\varepsilon}}^1$.

Corollary 5.5. Let $\alpha \geq 0$ and let $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n)z^n$, where $\widehat{f}(n) \geq 0$ for all nonnegative integers n and $\sum_{n=0}^{\infty} \widehat{f}(n)/(n+1) < \infty$. Then

$$Hf \in \mathcal{B}_{\log^{\alpha-1}}^1 \quad \text{if and only if} \quad \sum_{n=0}^{\infty} \widehat{f}(n) \frac{\log^\alpha(n+2)}{n+1} < \infty.$$

Moreover, there is a constant C independent of the function f , such that

$$\frac{1}{C} \|Hf\|_{\mathcal{B}_{\log^{\alpha-1}}^1} \leq \sum_{n=0}^{\infty} \widehat{f}(n) \frac{\log^\alpha(n+2)}{n+1} \leq C \|Hf\|_{\mathcal{B}_{\log^{\alpha-1}}^1}.$$

Proof. We have that $\widehat{Hf}(n) = \sum_{k=0}^{\infty} \widehat{f}(k)/(n+k+1) \downarrow 0$ as $n \rightarrow \infty$, because $\widehat{f}(k) \geq 0$ for all $k \geq 0$. Now the proof follows from the proof of part (b) of Theorem 5.3. \square

Corollary 5.6. Let $\alpha \geq 0$ and let $f(z) = \sum_{n=0}^{\infty} \widehat{f}(n)z^n \in \mathcal{H}(\mathbb{D})$, such that $\sum_{n=0}^{\infty} |\widehat{f}(n)| \log^\alpha(n+2)(n+1)^{-1} < \infty$. Then $Hf \in \mathcal{B}_{\log^{\alpha-1}}^1$.

Proof. Let $x_n = \operatorname{Re} \widehat{f}(n)$ and $y_n = \operatorname{Im} \widehat{f}(n)$ for all nonnegative integers n . Then, functions $g(z) = \sum_{n=0}^{\infty} x_n z^n$ and $h(z) = \sum_{n=0}^{\infty} y_n z^n$ are holomorphic in the unit disk \mathbb{D} . Now, let $x_n^+ = (|x_n| + x_n)/2$ and $x_n^- = (|x_n| - x_n)/2$ for all $n = 0, 1, \dots$. Then, $x_n^\pm \geq 0$, $x_n^\pm \leq |x_n| \leq |\widehat{f}(n)|$ and $x_n^+ - x_n^- = x_n$. Therefore, functions $g^+(z) = \sum_{n=0}^{\infty} x_n^+ z^n$ and $g^-(z) = \sum_{n=0}^{\infty} x_n^- z^n$ are holomorphic in the unit disk \mathbb{D} , and

$$\sum_{n=0}^{\infty} x_n^\pm \frac{\log^\alpha(n+2)}{n+1} \leq \sum_{n=0}^{\infty} |\widehat{f}(n)| \frac{\log^\alpha(n+2)}{n+1} < \infty.$$

Hence, by using Corollary 5.5, we find that

$$Hg^+ \in \mathcal{B}_{\log^{\alpha-1}}^1 \quad \text{and} \quad Hg^- \in \mathcal{B}_{\log^{\alpha-1}}^1.$$

Then, we have

$$\begin{aligned} \int_{\mathbb{D}} |(Hg)'(z)| \log^{\alpha-1} \frac{2}{1-|z|} dA(z) &= \int_{\mathbb{D}} |(Hg^+)'(z) - (Hg^-)'(z)| \log^{\alpha-1} \frac{2}{1-|z|} dA(z) \\ &\leq \int_{\mathbb{D}} (|(Hg^+)'(z)| + |(Hg^-)'(z)|) \log^{\alpha-1} \frac{2}{1-|z|} dA(z) < \infty, \end{aligned}$$

and we get $Hg \in \mathcal{B}_{\log^{\alpha-1}}^1$. In the same way, we prove that $Hh \in \mathcal{B}_{\log^{\alpha-1}}^1$. Then, we have that $Hf = Hg + iHh$, because of $f = g + ih$. Consequently,

$$\begin{aligned} \int_{\mathbb{D}} |(Hf)'(z)| \log^{\alpha-1} \frac{2}{1-|z|} dA(z) &= \int_{\mathbb{D}} |(Hg)'(z) + i(Hh)'(z)| \log^{\alpha-1} \frac{2}{1-|z|} dA(z) \\ &\leq \int_{\mathbb{D}} (|(Hg)'(z)| + |(Hh)'(z)|) \log^{\alpha-1} \frac{2}{1-|z|} dA(z) \\ &< \infty. \end{aligned}$$

Therefore, $Hf \in \mathcal{B}_{\log^{\alpha-1}}^1$. □

References

- [1] *M. Jevtić, B. Karapetrović*: Hilbert matrix operator on Besov spaces. *Publ. Math.* *90* (2017), 359–371. [MR](#) [doi](#)
- [2] *M. Jevtić, B. Karapetrović*: Libera operator on mixed norm spaces $H_{\nu}^{p,q,\alpha}$ when $0 < p < 1$. *Filomat* *31* (2017), 4641–4650. [MR](#)
- [3] *M. Jevtić, D. Vukotić, M. Arsenović*: Taylor Coefficients and Coefficient Multipliers of Hardy and Bergman-Type Spaces. *RSME Springer Series* 2, Springer, Cham, 2016. [zbl](#) [MR](#) [doi](#)
- [4] *B. Łanucha, M. Nowak, M. Pavlović*: Hilbert matrix operator on spaces of analytic functions. *Ann. Acad. Sci. Fenn., Math.* *37* (2012), 161–174. [zbl](#) [MR](#) [doi](#)
- [5] *M. Mateljević, M. Pavlović*: L^p -behaviour of the integral means of analytic functions. *Stud. Math.* *77* (1984), 219–237. [zbl](#) [MR](#) [doi](#)
- [6] *M. Pavlović*: Definition and properties of the libera operator on mixed norm spaces. *The Scientific World Journal* *2014* (2014), Article ID 590656, 15 pages. [doi](#)
- [7] *M. Pavlović*: Function Classes on the Unit Disc. An introduction. *De Gruyter Studies in Mathematics* 52, De Gruyter, Berlin, 2014. [zbl](#) [MR](#) [doi](#)
- [8] *M. Pavlović*: Logarithmic Bloch space and its predual. *Publ. Inst. Math. (Beograd) (N.S.)* *100(114)* (2016), 1–16. [zbl](#) [MR](#) [doi](#)
- [9] *K. Zhu*: Operator Theory in Function Spaces. *Monographs and Textbooks in Pure and Applied Mathematics* 139, Marcel Dekker, New York, 1990. [zbl](#) [MR](#)
- [10] *A. Zygmund*: Trigonometric Series. Vol. I, II. Cambridge University Press, Cambridge, 1959. [zbl](#) [MR](#)

Author's address: Boban Karapetrović, Faculty of Mathematics, University of Belgrade, Studentski trg 16, 11000 Belgrade, Serbia, e-mail: bkarapetrovic@matf.bg.ac.rs.